

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 6 : C07D 471/04, 401/06, 401/14, 519/00, A61K 31/415, 31/44 // (C07D 519/00, 498:00, 471:00) (C07D 519/00, 491:00, 471:00) (C07D 519/00, 471:00, 471:00)</p>	A1	<p>(11) International Publication Number: WO 99/31099</p> <p>(43) International Publication Date: 24 June 1999 (24.06.99)</p>																																													
<p>(21) International Application Number: PCT/US98/26568</p> <p>(22) International Filing Date: 14 December 1998 (14.12.98)</p> <p>(30) Priority Data:</p> <table border="0"> <tr><td>60/069,910</td><td>17 December 1997 (17.12.97)</td><td>US</td></tr> <tr><td>60/079,197</td><td>24 March 1998 (24.03.98)</td><td>US</td></tr> <tr><td>60/079,944</td><td>30 March 1998 (30.03.98)</td><td>US</td></tr> <tr><td>60/080,397</td><td>2 April 1998 (02.04.98)</td><td>US</td></tr> <tr><td>60/083,251</td><td>27 April 1998 (27.04.98)</td><td>US</td></tr> <tr><td>9810182.7</td><td>13 May 1998 (13.05.98)</td><td>GB</td></tr> <tr><td>9810882.2</td><td>20 May 1998 (20.05.98)</td><td>GB</td></tr> <tr><td>9810892.1</td><td>20 May 1998 (20.05.98)</td><td>GB</td></tr> <tr><td>9811283.2</td><td>26 May 1998 (26.05.98)</td><td>GB</td></tr> <tr><td>9812686.5</td><td>12 June 1998 (12.06.98)</td><td>GB</td></tr> <tr><td>60/092,588</td><td>13 July 1998 (13.07.98)</td><td>US</td></tr> <tr><td>60/092,624</td><td>13 July 1998 (13.07.98)</td><td>US</td></tr> <tr><td>60/099,948</td><td>11 September 1998 (11.09.98)</td><td>US</td></tr> <tr><td>9822331.6</td><td>13 October 1998 (13.10.98)</td><td>GB</td></tr> <tr><td>9822701.0</td><td>16 October 1998 (16.10.98)</td><td>GB</td></tr> </table> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p>		60/069,910	17 December 1997 (17.12.97)	US	60/079,197	24 March 1998 (24.03.98)	US	60/079,944	30 March 1998 (30.03.98)	US	60/080,397	2 April 1998 (02.04.98)	US	60/083,251	27 April 1998 (27.04.98)	US	9810182.7	13 May 1998 (13.05.98)	GB	9810882.2	20 May 1998 (20.05.98)	GB	9810892.1	20 May 1998 (20.05.98)	GB	9811283.2	26 May 1998 (26.05.98)	GB	9812686.5	12 June 1998 (12.06.98)	GB	60/092,588	13 July 1998 (13.07.98)	US	60/092,624	13 July 1998 (13.07.98)	US	60/099,948	11 September 1998 (11.09.98)	US	9822331.6	13 October 1998 (13.10.98)	GB	9822701.0	16 October 1998 (16.10.98)	GB	<p>(72) Inventors; and (75) Inventors/Applicants (for US only): DUGGAN, Mark, E. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). MEISSNER, Robert, S. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HUTCHINSON, John, H. [CA/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). HALCZENKO, Wasyl [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ASKEW, Ben, C. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). COLEMAN, Paul, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PATANE, Michael, A. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WANG, Jiabing [CN/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(81) Designated States: AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>
60/069,910	17 December 1997 (17.12.97)	US																																													
60/079,197	24 March 1998 (24.03.98)	US																																													
60/079,944	30 March 1998 (30.03.98)	US																																													
60/080,397	2 April 1998 (02.04.98)	US																																													
60/083,251	27 April 1998 (27.04.98)	US																																													
9810182.7	13 May 1998 (13.05.98)	GB																																													
9810882.2	20 May 1998 (20.05.98)	GB																																													
9810892.1	20 May 1998 (20.05.98)	GB																																													
9811283.2	26 May 1998 (26.05.98)	GB																																													
9812686.5	12 June 1998 (12.06.98)	GB																																													
60/092,588	13 July 1998 (13.07.98)	US																																													
60/092,624	13 July 1998 (13.07.98)	US																																													
60/099,948	11 September 1998 (11.09.98)	US																																													
9822331.6	13 October 1998 (13.10.98)	GB																																													
9822701.0	16 October 1998 (16.10.98)	GB																																													
<p>(54) Title: INTEGRIN RECEPTOR ANTAGONISTS</p> <p>(57) Abstract</p> <p>The present invention relates to compounds and derivatives thereof, their synthesis, and their use as integrin receptor antagonists. More particularly, the compounds of the present invention are antagonists of the integrin receptors $\alpha v\beta 3$, $\alpha v\beta 5$ and/or $\alpha v\beta 6$ and are useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, wound healing, viral disease, and tumor growth and metastasis.</p>																																															

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

TITLE OF THE INVENTION
INTEGRIN RECEPTOR ANTAGONISTS

5

CROSS-REFERENCE TO RELATED APPLICATIONS

The present invention is related to U.S. provisional applications Serial No. 60/069,910, filed December 17, 1997; 60/083,251, filed April 27, 1998; 60/092,588, filed July 13, 1998; 60/079,197, filed March
10 24, 1998; 60/079,944, filed March 30, 1998; 60/080,397, filed April 2, 1998; 60/092,624, filed July 13, 1998; and 60/099,948, filed September 11, 1998; the contents of each of which are hereby incorporated by reference.

FIELD OF THE INVENTION

15

The present invention relates to compounds and derivatives thereof, their synthesis, and their use as integrin receptor antagonists. More particularly, the compounds of the present invention are antagonists of the integrin receptors $\alpha v \beta 3$, $\alpha v \beta 5$, and/or $\alpha v \beta 6$ and are useful for inhibiting bone resorption, treating and preventing
20 osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, wound healing, viral disease, tumor growth, and metastasis.

BACKGROUND OF THE INVENTION

25

It is believed that a wide variety of disease states and conditions can be mediated by acting on integrin receptors and that integrin receptor antagonists represent a useful class of drugs. Integrin receptors are heterodimeric transmembrane proteins through which cells attach and communicate with extracellular matrices and other
30 cells (See S.B. Rodan and G.A. Rodan, "Integrin Function In Osteoclasts", *Journal of Endocrinology*, Vol. 154, S47- S56 (1997), which is incorporated by reference herein in its entirety).

In one aspect of the present invention, the compounds herein are useful for inhibiting bone resorption. Bone resorption is
35 mediated by the action of cells known as osteoclasts. Osteoclasts are

large multinucleated cells of up to about 400 μ m in diameter that resorb mineralized tissue, chiefly calcium carbonate and calcium phosphate, in vertebrates. Osteoclasts are actively motile cells that migrate along the surface of bone, and can bind to bone, secrete necessary acids and proteases, thereby causing the actual resorption of mineralized tissue from the bone. More specifically, osteoclasts are believed to exist in at least two physiological states, namely, the secretory state and the migratory or motile state. In the secretory state, osteoclasts are flat, attach to the bone matrix via a tight attachment zone (sealing zone), become highly polarized, form a ruffled border, and secrete lysosomal enzymes and protons to resorb bone. The adhesion of osteoclasts to bone surfaces is an important initial step in bone resorption. In the migratory or motile state, the osteoclasts migrate across bone matrix and do not take part in resorption until they again attach to bone.

Integrins are involved in osteoclast attachment, activation and migration. The most abundant integrin in osteoclasts, e.g., in rat, chicken, mouse and human osteoclasts, is an integrin receptor known as α v β 3, which is thought to interact in bone with matrix proteins that contain the RGD sequence. Antibodies to α v β 3 block bone resorption *in vitro* indicating that this integrin plays a key role in the resorptive process. There is increasing evidence to suggest that α v β 3 ligands can be used effectively to inhibit osteoclast mediated bone resorption *in vivo* in mammals.

The current major bone diseases of public concern are osteoporosis, hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, periarticular erosions in rheumatoid arthritis, Paget's disease, immobilization-induced osteopenia, and glucocorticoid-induced osteoporosis. All of these conditions are characterized by bone loss, resulting from an imbalance between bone resorption, i.e. breakdown, and bone formation, which continues throughout life at the rate of about 14% per year on the average. However, the rate of bone turnover differs from site to site; for example, it is higher in the trabecular bone of the vertebrae and the alveolar bone in the jaws than in the cortices of the long bones. The potential for bone loss is directly related to turnover and can amount to

over 5% per year in vertebrae immediately following menopause, a condition which leads to increased fracture risk.

In the United States, there are currently about 20 million people with detectable fractures of the vertebrae due to osteoporosis. In addition, there are about 250,000 hip fractures per year attributed to osteoporosis. This clinical situation is associated with a 12% mortality rate within the first two years, while 30% of the patients require nursing home care after the fracture.

Individuals suffering from all the conditions listed above would benefit from treatment with agents which inhibit bone resorption.

Additionally, $\alpha v \beta 3$ ligands have been found to be useful in treating and/or inhibiting restenosis, i.e. recurrence of stenosis after corrective surgery on the heart valve, atherosclerosis, diabetic retinopathy, macular degeneration, and angiogenesis, i.e. formation of new blood vessels. Moreover, it has been postulated that the growth of tumors depends on an adequate blood supply, which in turn is dependent on the growth of new vessels into the tumor; thus, inhibition of angiogenesis can cause tumor regression in animal models (See Harrison's Principles of Internal Medicine, 12th ed., 1991, which is incorporated by reference herein in its entirety). Therefore, $\alpha v \beta 3$ antagonists which inhibit angiogenesis can be useful in the treatment of cancer by inhibiting tumor growth (*See e.g., Brooks et al., Cell*, 79:1157-1164 (1994), which is incorporated by reference herein in its entirety).

Moreover, compounds of this invention can also inhibit neovascularization by acting as antagonists of the integrin receptor, $\alpha v \beta 5$. A monoclonal antibody for $\alpha v \beta 5$ has been shown to inhibit VEGF-induced angiogenesis in rabbit cornea and the chick chorioallantoic membrane model (*See M.C. Friedlander, et al., Science* 270, 1500-1502, (1995), which is incorporated by reference herein in its entirety). Thus, compounds that antagonize $\alpha v \beta 5$ are useful for treating and preventing macular degeneration, diabetic retinopathy, tumor growth, and metastasis.

Additionally, compounds of the instant invention can inhibit angiogenesis and inflammation by acting as antagonists of the integrin receptor, $\alpha v \beta 6$, which is expressed during the later stages of

wound healing and remains expressed until the wound is closed (See Christofidou-Solomidou, et al., "Expression and Function of Endothelial Cell α v Integrin Receptors in Wound-Induced Human Angiogenesis in Human Skin/SCID Mice Chimeras, American Journal of Pathology, Vol. 151, No. 4, pp. 975-983 (October 1997), which is incorporated by reference herein in its entirety). It is postulated that α v β 6 plays a role in the remodeling of the vasculature during the later stages of angiogenesis. Also, α v β 6 participates in the modulation of epithelial inflammation and is induced in response to local injury or inflammation (See Xiao-Zhu Huang, et al., "Inactivation of the Integrin β 6 Subunit Gene Reveals a Role of Epithelial Integrins in Regulating Inflammation in the Lungs and Skin," Journal of Cell Biology, Vol. 133, No.4, pp. 921-928 (May 1996), which is incorporated by reference herein in its entirety). Accordingly, compounds that antagonize α v β 6 are useful in treating or preventing cancer by inhibiting tumor growth and metastasis.

In addition, certain compounds of this invention antagonize both the α v β 3 and α v β 5 receptors. These compounds, referred to as "dual α v β 3/ α v β 5 antagonists," are useful for inhibiting bone resorption, treating and preventing osteoporosis, and inhibiting vascular restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, tumor growth, and metastasis.

In addition, certain compounds of this invention are useful as mixed α v β 3, α v β 5, and α v β 6 receptor antagonists.

It is therefore an object of the present invention to provide compounds which are useful as integrin receptor antagonists.

It is another object of the present invention to provide compounds which are useful as α v β 3 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as α v β 5 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as α v β 6 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as dual α v β 3/ α v β 5 receptor antagonists.

It is another object of the present invention to provide compounds which are useful as mixed $\alpha v\beta 3$, $\alpha v\beta 5$, and $\alpha v\beta 6$ receptor antagonists.

5 It is another object of the present invention to provide pharmaceutical compositions comprising integrin receptor antagonists.

It is another object of the present invention to provide methods for making the pharmaceutical compositions of the present invention.

10 It is another object of the present invention to provide methods for eliciting an integrin receptor antagonizing effect in a mammal in need thereof by administering the compounds and pharmaceutical compositions of the present invention.

15 It is another object of the present invention to provide compounds and pharmaceutical compositions useful for inhibiting bone resorption, restenosis, atherosclerosis, inflammation, viral disease, diabetic retinopathy, macular degeneration, angiogenesis, tumor growth, and metastasis.

20 It is another object of the present invention to provide compounds and pharmaceutical compositions useful for treating osteoporosis.




It is another object of the present invention to provide methods for inhibiting bone resorption, restenosis, atherosclerosis, inflammation, viral disease, diabetic retinopathy, macular degeneration, angiogenesis, tumor growth, and metastasis.

25 It is another object of the present invention to provide methods for treating osteoporosis.



These and other objects will become readily apparent from the detailed description which follows.



30 SUMMARY OF THE INVENTION

The present invention relates to compounds having a structural formula selected from the group consisting of




5





- 5 wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are substituted only with R¹⁰ and R¹²;

X is selected from the group consisting of



- 10 a 5- or 6-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R¹ substituent and the ring carbon atoms are unsubstituted or substituted with one or two R¹ substituents, and
- 15
- 20 a 9- to 14-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or

substituted with one R^1 substituent and the ring carbon atoms are unsubstituted or substituted with one or two R^1 substituents;

Y is selected from the group consisting of

- 5
- (CH₂)_m-,
 - (CH₂)_m-O-(CH₂)_n-,
 - (CH₂)_m-NR⁴-(CH₂)_n-,
 - (CH₂)_m-S-(CH₂)_n-,
 - 10 -(CH₂)_m-SO-(CH₂)_n-,
 - (CH₂)_m-SO₂-(CH₂)_n-,
 - (CH₂)_m-O-(CH₂)_n-O-(CH₂)_p-,
 - (CH₂)_m-O-(CH₂)_n-NR⁴-(CH₂)_p-,
 - (CH₂)_m-NR⁴-(CH₂)_n-NR⁴-(CH₂)_p-,
 - 15 -(CH₂)_m-O-(CH₂)_n-S-(CH₂)_p-,
 - (CH₂)_m-S-(CH₂)_n-S-(CH₂)_p-,
 - (CH₂)_m-NR⁴-(CH₂)_n-S-(CH₂)_p-,
 - (CH₂)_m-NR⁴-(CH₂)_n-O-(CH₂)_p-,
 - (CH₂)_m-S-(CH₂)_n-O-(CH₂)_p-,
 - 20 -(CH₂)_m-S-(CH₂)_n-NR⁴-(CH₂)_p-, and
 - (CH₂)_m-Z-(CH₂)_n-,

wherein Z is a 3- to 10-membered monocyclic or polycyclic aromatic or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R^1 substituent and the ring carbon atoms are unsubstituted or substituted with one or two R^1 substituents, and wherein any methylene (CH₂) carbon atom in Y, other than in R^4 , can be substituted by one or two R^3 substituents; and

30 wherein R^1 and R^2 are each independently selected from the group consisting of

hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₈ cycloalkyl,
C₃₋₈ cycloheteroalkyl, C₃₋₈ cycloalkyl C₁₋₆ alkyl,

C₃₋₈ cycloheteroalkyl C₁₋₆ alkyl, aryl, aryl C₁₋₈ alkyl, amino,
 amino C₁₋₈ alkyl, C₁₋₃ acylamino, C₁₋₃ acylamino C₁₋₈ alkyl,
 (C₁₋₆ alkyl)_pamino, (C₁₋₆ alkyl)_pamino C₁₋₈ alkyl,
 C₁₋₄ alkoxy, C₁₋₄ alkoxy C₁₋₆ alkyl, hydroxycarbonyl,
 5 hydroxycarbonyl C₁₋₆ alkyl, C₁₋₃ alkoxycarbonyl,
 C₁₋₃ alkoxycarbonyl C₁₋₆ alkyl, hydroxycarbonyl-
 C₁₋₆ alkyloxy, hydroxy, hydroxy C₁₋₆ alkyl, C₁₋₆ alkyloxy-
 C₁₋₆ alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,
 trifluoroethoxy, C₁₋₈ alkyl-S(O)_p, (C₁₋₈ alkyl)_paminocarbonyl,
 10 C₁₋₈ alkyloxycarbonylamino, (C₁₋₈ alkyl)_paminocarbonyloxy,
 (aryl C₁₋₈ alkyl)_pamino, (aryl)_pamino, aryl C₁₋₈
 alkylsulfonylamino, and C₁₋₈ alkylsulfonylamino;
 or two R¹ substituents, when on the same carbon atom, are taken
 together with the carbon atom to which they are attached to
 15 form a carbonyl group;

each R³ is independently selected from the group consisting of
 hydrogen,
 aryl,
 20 C₁₋₁₀ alkyl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 25 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 halogen,
 hydroxyl,
 oxo,
 30 trifluoromethyl,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,

C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 5 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 10 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 15 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 20 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 25 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 (aryl)_pamino,
 (aryl)_pamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino,
 30 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_paminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,

- arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
5 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
10 aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
15 aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)paminocarbonylamino,
(C₁₋₈ alkyl)paminocarbonylamino C₁₋₆ alkyl,
(aryl)paminocarbonylamino C₁₋₆ alkyl,
20 (aryl C₁₋₈ alkyl)paminocarbonylamino,
(aryl C₁₋₈ alkyl)paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)paminosulfonylamino,
(C₁₋₈ alkyl)paminosulfonylamino C₁₋₆ alkyl,
25 (aryl)paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)paminosulfonylamino,
(aryl C₁₋₈ alkyl)paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
30 arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
35 arylcarbonyl C₁₋₆ alkyl,

- aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
5 arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
(aryl)_paminocarbonyl C₁₋₆ alkyl,
10 (aryl C₁₋₈ alkyl)_paminocarbonyl, and
(aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl;
or two R³ substituents, when on the same carbon atom, are taken
together with the carbon atom to which they are attached to
form a carbonyl group or a cyclopropyl group,
15 wherein any of the alkyl groups of R³ are either unsubstituted or
substituted with one to three R¹ substituents,
and provided that each R³ is selected such that in the resultant
compound the carbon atom or atoms to which R³ is attached is itself
attached to no more than one heteroatom;
20 each R⁴ is independently selected from the group consisting of
hydrogen,
aryl,
aminocarbonyl,
25 C₃₋₈ cycloalkyl,
amino C₁₋₆ alkyl,
(aryl)_paminocarbonyl,
(aryl C₁₋₅ alkyl)_paminocarbonyl,
hydroxycarbonyl C₁₋₆ alkyl,
30 C₁₋₈ alkyl,
aryl C₁₋₆ alkyl,
(C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
(aryl C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
C₁₋₈ alkylsulfonyl,

- C₁₋₈ alkoxy carbonyl,
 aryloxy carbonyl,
 aryl C₁₋₈ alkoxy carbonyl,
 C₁₋₈ alkyl carbonyl,
 5 aryl carbonyl,
 aryl C₁₋₆ alkyl carbonyl,
 (C₁₋₈ alkyl)_p aminocarbonyl,
 aminosulfonyl,
 C₁₋₈ alkyl aminosulfonyl,
 10 (aryl)_p aminosulfonyl,
 (aryl C₁₋₈ alkyl)_p aminosulfonyl,
 aryl sulfonyl,
 aryl C₁₋₆ alkyl sulfonyl,
 C₁₋₆ alkyl thiocarbonyl,
 15 aryl thiocarbonyl, and
 aryl C₁₋₆ alkyl thiocarbonyl,
 wherein any of the alkyl groups of R⁴ are either unsubstituted or
 substituted with one to three R¹ substituents;
- 20 R⁵ and R⁶ are each independently selected from the group consisting of
 hydrogen,
 C₁₋₁₀ alkyl,
 aryl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 25 aryl-(CH₂)_r-S(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 30 halogen,
 hydroxyl,
 C₁₋₈ alkyl carbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxy carbonyl,

(C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 5 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 10 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 15 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 20 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₆ alkoxy,
 25 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 (aryl)_pamino,
 (aryl)_pamino C₁₋₆ alkyl,
 30 (aryl C₁₋₆ alkyl)_pamino,
 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_paminocarbonyloxy,

- C₁₋₈ alkylsulfonylamino,
arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
5 aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
10 aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
15 aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
20 (aryl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
25 (C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,
(aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
30 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,

C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 arylcarbonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 5 C₁₋₆ alkylthiocarbonylamino,
 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 arylthiocarbonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylthiocarbonylamino,
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 10 (C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminocarbonyl, and
 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl;
 or R⁵ and R⁶ are taken together with the carbon atom to which they are
 15 attached to form a carbonyl group,
 wherein any of the alkyl groups of R⁵ or R⁶ are either unsubstituted or
 substituted with one to three R¹ substituents,
 and provided that each R⁵ and R⁶ are selected such that in the resultant
 compound the carbon atom to which R⁵ and R⁶ are attached is itself
 20 attached to no more than one heteroatom;

R⁷ and R⁸ are each independently selected from the group consisting of
 hydrogen,
 C₁₋₁₀ alkyl,
 25 aryl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 30 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 halogen,
 hydroxyl,
 C₁₋₈ alkylcarbonylamino,

aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 5 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 10 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 15 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 20 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 25 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 30 (aryl)_pamino,
 (aryl)_pamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino,
 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,

aryl C₁₋₆ alkylcarbonyloxy,
(C₁₋₆ alkyl)_paminocarbonyloxy,
C₁₋₈ alkylsulfonylamino,
arylcarbonylamino,
5 arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
10 C₁₋₈ alkoxycarbonylamino,
C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
15 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
20 (C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl)_paminocarbonylamino C₁₋₆ alkyl,
arylaminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
25 (aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
30 (aryl C₁₋₈ alkyl)_paminosulfonylamino,
(aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,

aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
5 arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
10 arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
(aryl)_paminocarbonyl C₁₋₆ alkyl,
15 (aryl C₁₋₈ alkyl)_paminocarbonyl,
(aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl, and
C₇₋₂₀ polycyclyl C₀₋₈ alkylsulfonylamino;

wherein any of the alkyl groups of R⁷ and R⁸ are either unsubstituted or substituted with one to three R¹ substituents,
20 and provided that each R⁷ and R⁸ are selected such that in the resultant compound the carbon atom to which R⁷ and R⁸ are attached is itself attached to no more than one heteroatom;

R⁹ is selected from the group consisting of
25 hydrogen,
C₁₋₈ alkyl,
aryl,
aryl C₁₋₈ alkyl,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
30 aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
C₁₋₈ alkylaminocarbonylmethylene, and
C₁₋₈ dialkylaminocarbonylmethylene;

R10, R11, R12 and R13 are each independently selected from the group consisting of

- hydrogen,
- C₁₋₈ alkyl,
- 5 aryl,
- halogen,
- hydroxyl,
- aminocarbonyl,
- C₃₋₈ cycloalkyl,
- 10 amino C₁₋₆ alkyl,
- (aryl)_paminocarbonyl,
- hydroxycarbonyl,
- (aryl C₁₋₅ alkyl)_paminocarbonyl,
- hydroxycarbonyl C₁₋₆ alkyl,
- 15 aryl C₁₋₆ alkyl,
- (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
- (aryl C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
- C₁₋₈ alkylsulfonyl,
- C₁₋₈ alkoxycarbonyl,
- 20 aryloxycarbonyl,
- aryl C₁₋₈ alkoxycarbonyl,
- C₁₋₈ alkylcarbonyl,
- arylcarbonyl,
- aryl C₁₋₆ alkylcarbonyl,
- 25 (C₁₋₈ alkyl)_paminocarbonyl,
- aminosulfonyl,
- C₁₋₈ alkylaminosulfonyl,
- (aryl)_paminosulfonyl,
- (aryl C₁₋₈ alkyl)_paminosulfonyl,
- 30 C₁₋₆ alkylsulfonyl,
- arylsulfonyl,
- aryl C₁₋₆ alkylsulfonyl,
- aryl C₁₋₆ alkylcarbonyl,
- C₁₋₆ alkylthiocarbonyl,

- arylthiocarbonyl,
 aryl C₁₋₆ alkylthiocarbonyl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 5 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 HC≡C-(CH₂)_t-,
 10 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 15 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 20 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 25 C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_pamino,
 aminocarbonyl C₁₋₆ alkyl,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 30 (aryl)_pamino,
 (aryl)_pamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino,
 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 35 aryl C₁₋₆ alkylcarbonyloxy,

- (C₁₋₆ alkyl)paminocarbonyloxy,
C₁₋₈ alkylsulfonylamino,
arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
5 arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
10 aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
15 arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)paminocarbonylamino,
20 (C₁₋₈ alkyl)paminocarbonylamino C₁₋₆ alkyl,
(aryl)paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)paminocarbonylamino,
(aryl C₁₋₈ alkyl)paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
25 (C₁₋₈ alkyl)paminosulfonylamino,
(C₁₋₈ alkyl)paminosulfonylamino C₁₋₆ alkyl,
(aryl)paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)paminosulfonylamino,
(aryl C₁₋₈ alkyl)paminosulfonylamino C₁₋₆ alkyl,
30 C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,

- C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
5 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
10 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
(aryl)_paminocarbonyl C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonyl, and
(aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl; or
15 R¹⁰ and R¹² are taken together with the carbon atoms to which
they are attached to form a 5- to 7-membered monocyclic aromatic
or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms
selected from the group consisting of N, O, and S wherein the ring
nitrogen atoms are unsubstituted or substituted with one R¹
20 substituent and the ring carbon atoms are unsubstituted or
substituted with one or two R¹ substituents,
and wherein any of the alkyl groups of R¹⁰, R¹¹, R¹², and R¹³ are either
unsubstituted or substituted with one to three R¹ substituents;
- 25 wherein
each m is independently an integer from 0 to 6;
each n is independently an integer from 0 to 6
each p is independently an integer from 0 to 2;
each r is independently an integer from 1 to 3;
30 each s is independently an integer from 0 to 3;
each t is independently an integer from 0 to 3; and
each v is independently an integer from 0 to 2;

and the pharmaceutically acceptable salts thereof.

The present invention also relates to pharmaceutical compositions comprising the compounds of the present invention and a pharmaceutically acceptable carrier.

5 The present invention also relates to methods for making the pharmaceutical compositions of the present invention.


The present invention also relates to methods for eliciting an integrin receptor antagonizing effect in a mammal in need thereof by administering the compounds and pharmaceutical compositions of the present invention.

10 The present invention also relates to methods for inhibiting bone resorption, restenosis, atherosclerosis, inflammation, viral disease, diabetic retinopathy, macular degeneration, angiogenesis, wound healing, tumor growth, and metastasis by administering the compounds and pharmaceutical compositions of the present invention.




15 The present invention also relates to methods for treating osteoporosis by administering the compounds and pharmaceutical compositions of the present invention.

DETAILED DESCRIPTION OF THE INVENTION



20 The present invention relates to compounds useful as integrin receptor antagonists. Compounds of the present invention are described by the following structural formulas selected from the group consisting of



25




5




, and

10



wherein the dotted line a represents a single or a double bond, provided
that when a represents a double bond, the double bond carbon atoms are
5 substituted only with R¹⁰ and R¹²;

X is selected from the group consisting of



10 a 5- or 6-membered monocyclic aromatic or nonaromatic ring
system having 0, 1, 2, 3 or 4 heteroatoms selected from the group
consisting of N, O, and S wherein the ring nitrogen atoms are
unsubstituted or substituted with one R¹ substituent and the ring
carbon atoms are unsubstituted or substituted with one or two R¹
substituents, and

15 a 9- to 14-membered polycyclic ring system, wherein one or more
of the rings is aromatic, and wherein the polycyclic ring system
has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of
N, O, and S wherein the ring nitrogen atoms are unsubstituted or
substituted with one R¹ substituent and the ring carbon atoms are
20 unsubstituted or substituted with one or two R¹ substituents;

Y is selected from the group consisting of

25 -(CH₂)_m-,
-(CH₂)_m-O-(CH₂)_n-,
-(CH₂)_m-NR⁴-(CH₂)_n-,

5 $-(CH_2)_m-S-(CH_2)_n-$,
 $-(CH_2)_m-SO-(CH_2)_n-$,
 $-(CH_2)_m-SO_2-(CH_2)_n-$,
 $-(CH_2)_m-O-(CH_2)_n-O-(CH_2)_p-$,
 $-(CH_2)_m-O-(CH_2)_n-NR^4-(CH_2)_p-$,
 $-(CH_2)_m-NR^4-(CH_2)_n-NR^4-(CH_2)_p-$,
 $-(CH_2)_m-O-(CH_2)_n-S-(CH_2)_p-$,
 $-(CH_2)_m-S-(CH_2)_n-S-(CH_2)_p-$,
 $-(CH_2)_m-NR^4-(CH_2)_n-S-(CH_2)_p-$,
 10 $-(CH_2)_m-NR^4-(CH_2)_n-O-(CH_2)_p-$,
 $-(CH_2)_m-S-(CH_2)_n-O-(CH_2)_p-$,
 $-(CH_2)_m-S-(CH_2)_n-NR^4-(CH_2)_p-$, and
 $-(CH_2)_m-Z-(CH_2)_n-$,

15 wherein Z is a 3- to 10-membered monocyclic or polycyclic aromatic or
 nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected
 from the group consisting of N, O, and S wherein the 3- to 10-membered
 monocyclic or polycyclic aromatic or nonaromatic ring system is either
 unsubstituted or substituted with one or two R^1 substituents, and
 wherein any methylene (CH_2) carbon atom in Y, other than in R^4 , can
 20 be substituted by one or two R^3 substituents; and

 wherein R^1 and R^2 are each independently selected from the group
 consisting of

25 hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₈ cycloalkyl,
 C₃₋₈ cycloheteroalkyl, C₃₋₈ cycloalkyl C₁₋₆ alkyl,
 C₃₋₈ cycloheteroalkyl C₁₋₆ alkyl, aryl, aryl C₁₋₈ alkyl, amino,
 amino C₁₋₈ alkyl, C₁₋₃ acylamino, C₁₋₃ acylamino C₁₋₈ alkyl,
 (C₁₋₆ alkyl)_pamino, (C₁₋₆ alkyl)_pamino C₁₋₈ alkyl,
 C₁₋₄ alkoxy, C₁₋₄ alkoxy C₁₋₆ alkyl, hydroxycarbonyl,
 30 hydroxycarbonyl C₁₋₆ alkyl, C₁₋₃ alkoxycarbonyl,
 C₁₋₃ alkoxycarbonyl C₁₋₆ alkyl, hydroxycarbonyl-
 C₁₋₆ alkyloxy, hydroxy, hydroxy C₁₋₆ alkyl, C₁₋₆ alkyloxy-
 C₁₋₆ alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,
 trifluoroethoxy, C₁₋₈ alkyl-S(O)_p, (C₁₋₈ alkyl)_paminocarbonyl,

C₁₋₈ alkyloxycarbonylamino, (C₁₋₈ alkyl)_paminocarbonyloxy,
 (aryl C₁₋₈ alkyl)_pamino, (aryl)_pamino, aryl C₁₋₈
 alkylsulfonylamino, and C₁₋₈ alkylsulfonylamino;
 or two R¹ substituents, when on the same carbon atom, are taken
 5 together to form a carbonyl group;

each R³ is independently selected from the group consisting of
 hydrogen,
 aryl,
 10 C₁₋₁₀ alkyl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 15 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 halogen,
 hydroxyl,
 oxo,
 20 trifluoromethyl,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 25 C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 30 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,

- $\text{HC}\equiv\text{C}-(\text{CH}_2)_t-$,
 $\text{C}_{1-6} \text{ alkyl}-\text{C}\equiv\text{C}-(\text{CH}_2)_t-$,
 $\text{C}_{3-7} \text{ cycloalkyl}-\text{C}\equiv\text{C}-(\text{CH}_2)_t-$,
 $\text{aryl}-\text{C}\equiv\text{C}-(\text{CH}_2)_t-$,
5 $\text{C}_{1-6} \text{ alkylaryl}-\text{C}\equiv\text{C}-(\text{CH}_2)_t-$,
 $\text{CH}_2=\text{CH}-(\text{CH}_2)_t-$,
 $\text{C}_{1-6} \text{ alkyl}-\text{CH}=\text{CH}-(\text{CH}_2)_t-$,
 $\text{C}_{3-7} \text{ cycloalkyl}-\text{CH}=\text{CH}-(\text{CH}_2)_t-$,
 $\text{aryl}-\text{CH}=\text{CH}-(\text{CH}_2)_t-$,
10 $\text{C}_{1-6} \text{ alkylaryl}-\text{CH}=\text{CH}-(\text{CH}_2)_t-$,
 $\text{C}_{1-6} \text{ alkyl}-\text{SO}_2-(\text{CH}_2)_t-$,
 $\text{C}_{1-6} \text{ alkylaryl}-\text{SO}_2-(\text{CH}_2)_t-$,
 $\text{C}_{1-6} \text{ alkoxy}$,
 $\text{aryl C}_{1-6} \text{ alkoxy}$,
15 $\text{aryl C}_{1-6} \text{ alkyl}$,
 $(\text{C}_{1-6} \text{ alkyl})_{\text{pamino C}_{1-6} \text{ alkyl}}$,
 $(\text{aryl})_{\text{pamino}}$,
 $(\text{aryl})_{\text{pamino C}_{1-6} \text{ alkyl}}$,
 $(\text{aryl C}_{1-6} \text{ alkyl})_{\text{pamino}}$,
20 $(\text{aryl C}_{1-6} \text{ alkyl})_{\text{pamino C}_{1-6} \text{ alkyl}}$,
 arylcarbonyloxy ,
 $\text{aryl C}_{1-6} \text{ alkylcarbonyloxy}$,
 $(\text{C}_{1-6} \text{ alkyl})_{\text{paminocarbonyloxy}}$,
 $\text{C}_{1-8} \text{ alkylsulfonylamino}$,
25 arylsulfonylamino ,
 $\text{C}_{1-8} \text{ alkylsulfonylamino C}_{1-6} \text{ alkyl}$,
 $\text{arylsulfonylamino C}_{1-6} \text{ alkyl}$,
 $\text{aryl C}_{1-6} \text{ alkylsulfonylamino}$,
 $\text{aryl C}_{1-6} \text{ alkylsulfonylamino C}_{1-6} \text{ alkyl}$,
30 $\text{C}_{1-8} \text{ alkoxycarbonylamino}$,
 $\text{C}_{1-8} \text{ alkoxycarbonylamino C}_{1-8} \text{ alkyl}$,
 $\text{aryloxycarbonylamino C}_{1-8} \text{ alkyl}$,
 $\text{aryl C}_{1-8} \text{ alkoxycarbonylamino}$,
 $\text{aryl C}_{1-8} \text{ alkoxycarbonylamino C}_{1-8} \text{ alkyl}$,

- C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcabonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
5 aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl)_paminocarbonylamino C₁₋₆ alkyl,
10 (aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
15 (aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,
(aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
20 arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
25 arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
30 arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
(aryl)_paminocarbonyl C₁₋₆ alkyl,

(aryl C₁₋₈ alkyl)_paminocarbonyl, and
(aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl;
or two R³ substituents, when on the same carbon atom, are taken
together with the carbon atom to which they are attached to
5 form a carbonyl group or a cyclopropyl group,
wherein any of the alkyl groups of R³ are either unsubstituted or
substituted with one to three R¹ substituents,
and provided that each R³ is selected such that in the resultant
compound the carbon atom or atoms to which R³ is attached is itself
10 attached to no more than one heteroatom;

each R⁴ is independently selected from the group consisting of
hydrogen;
aryl,
15 aminocarbonyl,
C₃₋₈ cycloalkyl,
amino C₁₋₆ alkyl,
(aryl)_paminocarbonyl,
(aryl C₁₋₅ alkyl)_paminocarbonyl,
20 hydroxycarbonyl C₁₋₆ alkyl,
C₁₋₈ alkyl,
aryl C₁₋₆ alkyl,
(C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
(aryl C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
25 C₁₋₈ alkylsulfonyl,
C₁₋₈ alkoxy carbonyl,
aryloxy carbonyl,
aryl C₁₋₈ alkoxy carbonyl,
C₁₋₈ alkyl carbonyl,
30 aryl carbonyl,
aryl C₁₋₆ alkyl carbonyl,
(C₁₋₈ alkyl)_paminocarbonyl,
aminosulfonyl,
C₁₋₈ alkylaminosulfonyl,

(aryl)_paminosulfonyl,
 (aryl C₁₋₈ alkyl)_paminosulfonyl,
 arylsulfonyl,
 arylC₁₋₆ alkylsulfonyl,
 5 C₁₋₆ alkylthiocarbonyl,
 arylthiocarbonyl, and
 aryl C₁₋₆ alkylthiocarbonyl,
 wherein any of the alkyl groups of R⁴ are either unsubstituted or
 substituted with one to three R¹ substituents;

10 R⁵ and R⁶ are each independently selected from the group consisting of
 hydrogen,
 C₁₋₁₀ alkyl,
 aryl,
 15 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 20 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 halogen,
 hydroxyl,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 25 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 30 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,

- hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 5 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 10 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 15 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 (aryl)_pamino,
 20 (aryl)_pamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino,
 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 25 (C₁₋₆ alkyl)_paminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,
 arylsulfonylamino,
 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
 arylsulfonylamino C₁₋₆ alkyl,
 30 aryl C₁₋₆ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
 C₁₋₈ alkoxycarbonylamino,
 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 aryloxycarbonylamino C₁₋₈ alkyl,
 35 aryl C₁₋₈ alkoxycarbonylamino,

- aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcabonylamino C₁₋₆ alkyl,
5 aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
10 (aryl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
15 (C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,
(aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
20 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
25 C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcabonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
30 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,

(aryl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminocarbonyl, and
 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl;

or R⁵ and R⁶ are taken together with the carbon atom to which
 5 they are attached to form a carbonyl group,
 wherein any of the alkyl groups of R⁵ or R⁶ are either unsubstituted or
 substituted with one to three R¹ substituents,
 and provided that each R⁵ and R⁶ are selected such that in the resultant
 compound the carbon atom to which R⁵ and R⁶ are attached is itself
 10 attached to no more than one heteroatom;

R⁷ and R⁸ are each independently selected from the group consisting of
 hydrogen,
 C₁₋₁₀ alkyl,
 15 aryl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 20 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 halogen,
 hydroxyl,
 C₁₋₈ alkylcarbonylamino,
 25 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 30 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,

- aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 5 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 10 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 15 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 20 (aryl)_pamino,
 (aryl)_pamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino,
 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 25 aryl C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_paminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,
 arylsulfonylamino,
 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
 30 arylsulfonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
 C₁₋₈ alkoxycarbonylamino,
 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,

aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
5 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
10 (C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
15 aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,
20 (aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
25 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
30 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,

aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
(aryl)_paminocarbonyl C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonyl,
5 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl, and
C₇₋₂₀ polycyclyl C₀₋₈ alkylsulfonylamino;

wherein any of the alkyl groups of R⁷ and R⁸ are either unsubstituted or substituted with one to three R¹ substituents,
and provided that each R⁷ and R⁸ are selected such that in the resultant
10 compound the carbon atom to which R⁷ and R⁸ are attached is itself
attached to no more than one heteroatom;

R⁹ is selected from the group consisting of
hydrogen,
15 C₁₋₈ alkyl,
aryl,
aryl C₁₋₈ alkyl,
C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
20 C₁₋₈ alkylaminocarbonylmethylene, and
C₁₋₈ dialkylaminocarbonylmethylene;

R¹⁰, R¹¹, R¹² and R¹³ are each independently selected from the group
consisting of
25 hydrogen,
C₁₋₈ alkyl,
aryl,
halogen,
hydroxyl,
30 aminocarbonyl,
C₃₋₈ cycloalkyl,
amino C₁₋₆ alkyl,
(aryl)_paminocarbonyl,
hydroxycarbonyl,

- (aryl C₁₋₅ alkyl)_paminocarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 5 (aryl C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
 C₁₋₈ alkylsulfonyl,
 C₁₋₈ alkoxycarbonyl,
 aryloxycarbonyl,
 aryl C₁₋₈ alkoxycarbonyl,
 10 C₁₋₈ alkylcarbonyl,
 arylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 aminosulfonyl,
 15 C₁₋₈ alkylaminosulfonyl,
 (aryl)_paminosulfonyl,
 (aryl C₁₋₈ alkyl)_paminosulfonyl,
 C₁₋₆ alkylsulfonyl,
 arylsulfonyl,
 20 aryl C₁₋₆ alkylsulfonyl,
 aryl C₁₋₆ alkylcarbonyl,
 C₁₋₆ alkylthiocarbonyl,
 arylthiocarbonyl,
 aryl C₁₋₆ alkylthiocarbonyl,
 25 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 30 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,

- C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 5 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₈ alkylcarbonylamino,
 10 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_pamino,
 15 aminocarbonyl C₁₋₆ alkyl,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 (aryl)_pamino,
 (aryl)_pamino C₁₋₆ alkyl,
 20 (aryl C₁₋₆ alkyl)_pamino,
 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_paminocarbonyloxy,
 25 C₁₋₈ alkylsulfonylamino,
 arylsulfonylamino,
 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
 arylsulfonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylsulfonylamino,
 30 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
 C₁₋₈ alkoxycarbonylamino,
 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 aryloxycarbonylamino C₁₋₈ alkyl,
 aryl C₁₋₈ alkoxycarbonylamino,

- aryl C₁₋₈ alkoxy-carbonylamino C₁₋₈ alkyl,
C₁₋₈ alkyl-carbonylamino,
C₁₋₈ alkyl-carbonylamino C₁₋₆ alkyl,
aryl-carbonylamino C₁₋₆ alkyl,
5 aryl C₁₋₆ alkyl-carbonylamino,
aryl C₁₋₆ alkyl-carbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
10 (aryl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
15 (C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,
(aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylsulfonyl,
20 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
25 C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
30 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,


(aryl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminocarbonyl, and
 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl; or
 R¹⁰ and R¹² are taken together with the carbon atoms to which
 5 they are attached to form a 5- to 7-membered monocyclic aromatic
 or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms
 selected from the group consisting of N, O, and S wherein the ring
 nitrogen atoms are unsubstituted or substituted with one R¹
 10 substituent and the ring carbon atoms are unsubstituted or
 substituted with one or two R¹ substituents,
 and wherein any of the alkyl groups of R¹⁰, R¹¹, R¹², and R¹³ are either
 unsubstituted or substituted with one to three R¹ substituents;

wherein


15 each m is independently an integer from 0 to 6;
 each n is independently an integer from 0 to 6
 each p is independently an integer from 0 to 2;
 each r is independently an integer from 1 to 3;
 each s is independently an integer from 0 to 3;
 20 each t is independently an integer from 0 to 3; and
 each v is independently an integer from 0 to 2;

and the pharmaceutically acceptable salts thereof.

25 In one embodiment of the present invention, compounds are
 described by the following structural formulas selected from the group
 consisting of




30



wherein the dotted line a represents a single or a double bond, provided
 5 that when a represents a double bond, the double bond carbon atoms are
 substituted only with R¹⁰ and R¹².


In a class of this embodiment of the present invention,
 compounds are described by the following structural formula

10



wherein the dotted line a represents a single or a double bond, provided
 that when a represents a double bond, the double bond carbon atoms are
 15 substituted only with R¹⁰ and R¹².

In a subclass of this class of the present invention,
 compounds are described by the following structural formula




In the compounds of the present invention, X is preferably
a 6-membered monocyclic aromatic ring system having 1 or 2 nitrogen
5 atoms wherein each ring carbon atom is unsubstituted or substituted
with one R¹ substituent, or

a 9- to 14-membered polycyclic ring system, wherein one or more of the
rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3
10 or 4 heteroatoms selected from the group consisting of N, O, and S
wherein the ring nitrogen atoms are unsubstituted or substituted with
one R¹ substituent and the ring carbon atoms are unsubstituted or
substituted with one or two R¹ substituents.


15

More preferably, X is selected from the group consisting of



20

Most preferably X is




In the compounds of the present invention, Y is preferably
5 selected from the group consisting of

- (CH₂)_m-,
- (CH₂)_m-O-(CH₂)_n-,
- (CH₂)_m-NR⁴-(CH₂)_n-,
- 10 -(CH₂)_m-S-(CH₂)_n-,
- (CH₂)_m-SO-(CH₂)_n-,
- (CH₂)_m-SO₂-(CH₂)_n-,
- (CH₂)_m-O-(CH₂)_n-O-(CH₂)_p-,
- (CH₂)_m-O-(CH₂)_n-NR⁴-(CH₂)_p-,
- 15 -(CH₂)_m-NR⁴-(CH₂)_n-NR⁴-(CH₂)_p-, and
- (CH₂)_m-NR⁴-(CH₂)_n-O-(CH₂)_p-,

wherein any methylene (CH₂) carbon atom in Y, other than in R⁴, can be substituted by one or two R³ substituents.

20 More preferably Y is selected from the group consisting of



wherein any methylene (CH₂) carbon atom in Y, other than in R⁴, can
25 be substituted by one or two R³ substituents.

Most preferably Y is (CH₂)_m or (CH₂)_m-NR⁴-(CH₂)_n
wherein any methylene (CH₂) carbon atom in Y, other than R⁴, can be
substituted by one or two R³ substituents.

30

In the compounds of the present invention, R¹ and R² are preferably selected from the group consisting of hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, C₃-8 cycloheteroalkyl, hydroxy, nitro, cyano, trifluoromethyl, and trifluoromethoxy.

More preferably, R¹ and R² are selected from the group consisting of hydrogen, halogen, C₁-10 alkyl, C₃-8 cycloalkyl, trifluoromethyl, and trifluoromethoxy.

In the compounds of the present invention, R³ is preferably selected from the group consisting of

hydrogen,
fluoro,
trifluoromethyl,
aryl,
C₁-8 alkyl,
arylC₁-6 alkyl
hydroxyl,
oxo,
arylamino-carbonyl,
aryl C₁-5 alkylaminocarbonyl,
aminocarbonyl, and
aminocarbonyl C₁-6 alkyl.

More preferably, R³ is selected from the group consisting of
fluoro,
aryl,
C₁-8 alkyl,
arylC₁-6 alkyl
hydroxyl,
oxo, and
arylamino-carbonyl.

In the compounds of the present invention, R⁴ is preferably selected from the group consisting of

hydrogen,
aryl,
C3-8 cycloalkyl,
C1-8 alkyl,
5 C1-8 alkylcarbonyl,
arylcarbonyl,
C1-6 alkylsulfonyl,
arylsulfonyl,
arylC1-6alkylsulfonyl,
10 arylC1-6alkylcarbonyl,
C1-8alkylaminocarbonyl,
arylC1-5alkylaminocarbonyl,
arylC1-8alkoxycarbonyl, and
C1-8alkoxycarbonyl.

15

More preferably, R^4 is selected from the group consisting of

hydrogen,
C1-8 alkyl,
C1-8 alkylcarbonyl,
20 arylcarbonyl,
arylC1-6alkylcarbonyl,
C1-6 alkylsulfonyl,
arylsulfonyl, and
arylC1-6alkylsulfonyl.

25

In one embodiment of the present invention, R^5 and R^6 are each independently selected from the group consisting of

hydrogen,
aryl,
30 C1-8 alkyl,
aryl-C \equiv C-(CH₂)_t-,
aryl C1-6 alkyl,
CH₂=CH-(CH₂)_t-, and
HC \equiv C-(CH₂)_t-.

In a class of this embodiment of the present invention, R^6 is hydrogen and R^5 is selected from the group consisting of

hydrogen,
aryl,
5 C₁₋₈ alkyl,
aryl-C \equiv C-(CH₂)_t-,
aryl C₁₋₆ alkyl,
CH₂=CH-(CH₂)_t-, and
HC \equiv C-(CH₂)_t-.

10 In a subclass of this class of the present invention, R^6 , R^7 , and R^8 are each hydrogen and R^5 is selected from the group consisting of

hydrogen,
aryl,
15 C₁₋₈ alkyl,
aryl-C \equiv C-(CH₂)_t-,
aryl C₁₋₆ alkyl,
CH₂=CH-(CH₂)_t-, and
HC \equiv C-(CH₂)_t-.

20 In another embodiment of the present invention, R^7 and R^8 are each independently selected from the group consisting of

hydrogen,
aryl,
C₁₋₈ alkylcarbonylamino,
25 arylcarbonylamino,
C₁₋₈ alkylsulfonylamino,
arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
30 aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxy carbonylamino,
C₁₋₈ alkoxy carbonylamino C₁₋₈ alkyl,
aryloxy carbonylamino C₁₋₈ alkyl,

aryl C₁₋₈ alkoxycarbonylamino,
 aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
 arylcarbonylamino C₁₋₆ alkyl,
 5 aryl C₁₋₆ alkylcarbonylamino,
 aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
 aminocarbonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)_paminocarbonylamino,
 (C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
 10 (aryl)_paminocarbonylamino C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminocarbonylamino,
 (aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
 aminosulfonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)_paminosulfonylamino,
 15 (C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
 (aryl)_paminosulfonylamino C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminosulfonylamino,
 (aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
 C₁₋₆ alkylthiocarbonylamino,
 20 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 arylthiocarbonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylthiocarbonylamino, and
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl.

25 In a class of this embodiment of the present invention, R⁸ is
 hydrogen and R⁷ is selected from the group consisting of consisting of
 hydrogen,
 aryl,
 C₁₋₈ alkylcarbonylamino,
 30 aryl C₁₋₆ alkylcarbonylamino,
 arylcarbonylamino,
 C₁₋₈ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino,
 arylsulfonylamino,

C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino,
arylamino carbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino,
5 (aryl C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino, and
(aryl C₁₋₈ alkyl)_paminosulfonylamino.

In a subclass of this class of the present invention, R⁵, R⁶,
and R⁸ are each hydrogen and R⁷ is selected from the group consisting
10 of

hydrogen,
aryl,
C₁₋₈ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino,
15 arylcarbonylamino,
C₁₋₈ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino,
arylsulfonylamino,
C₁₋₈ alkoxycarbonylamino,
20 aryl C₁₋₈ alkoxycarbonylamino,
arylamino carbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino, and
25 (aryl C₁₋₈ alkyl)_paminosulfonylamino.

In the compounds of the present invention, R⁹ is preferably
selected from the group consisting of hydrogen, methyl, and ethyl.

More preferably, R⁹ is hydrogen.

30 In the compounds of the present invention, R¹⁰, R¹¹, R¹²,
and R¹³ are preferably each independently selected from the group
consisting of hydrogen, aryl, C₁₋₆ alkyl, and arylC₁₋₆ alkyl.

In the compounds of the present invention, m is preferably
an integer from 0 to 4, and more preferably from 0 to 3.

In the compounds of the present invention, n is preferably an integer from 0 to 4, more preferably from 0 to 3.



In the compounds of the present invention, r is preferably an integer from 1 to 2.

5 In the compounds of the present invention, s is preferably an integer from 0 to 2.

In the compounds of the present invention, t is preferably an integer from 0 to 2, more preferably from 0 to 1.


In the compounds of the present invention, v is preferably 0.

10 In certain embodiments of the present invention, the compounds correspond to the formulas with the following designated stereochemistry at the carbon atom where R⁵ and R⁶ are attached:



15

, and



;

wherein the substituents X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², and R¹³ and the subscripts a, m, n, p, r, s, t, and v are as described above.

5 Illustrative but nonlimiting examples of compounds of the present invention that are useful as integrin receptor antagonists are the following:

- 10 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 15 3-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 20 3(R)-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3(S)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 30 3-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

5 3(R)-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

10 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15 3(R)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,

3(R)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,

25 3-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,

30 3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
trifluoroacetate salt,
- 3(S)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
bis(trifluoroacetate) salt,
- 3(R)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
25 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-
35 propionic acid,

3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

5

3-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

10 3-(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(R)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15

3-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20 3-(S)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(R)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25 3-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

30

3-(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

35 3-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3(S)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 2(S)-Benzenesulfonylamino-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
30 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-
imidazolidin-1-yl}-pent-4-enoic acid,

- 3(R)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 5 3-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 10 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 15 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(R)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 20 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 25 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 35 3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

- 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5
- 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10
- 3-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15
- 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20
- 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25
- 3-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30
- 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

3-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

- 5 3-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

10

3(R)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 15 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 20 3-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

25

3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 30 3-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

5 3-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

10 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

15 3-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

20 3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

3-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

25 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

30 3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

35 3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

5 3-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

10 Further illustrative of the present invention are the compounds selected from the group consisting of

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15

3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20 3(S)-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(3-Fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25 3(S)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

30

3(S)-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

35 3(R)-(Ethynyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 10 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- 15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(S)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- bis(trifluoroacetate) salt,
- 30 3(R)-(4-Methoxyquinolin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
- [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(R)-(6-Amino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(S)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3-(R)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3-(S)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3-(R)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35 3-(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3-(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35 3-(S)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35 3-(R)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 5 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 10 3(R)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-
15 imidazolidin-1-yl)-pent-4-enoic acid,
- 3(R)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-
imidazolidin-1-yl)-pent-4-enoic acid,
- 20 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-
yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-
yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 25 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-
ethoxy-pyridin-3-yl)-propionic acid,
- 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(R)-(5-
30 ethoxy-pyridin-3-yl)-propionic acid,
- 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-
yl)propionic acid,
- 35

- 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 5 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 10 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 15 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 20 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 25 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 30 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,
- 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,
- 3(R)-(Furo[2,3-b]pyridin-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl}propionic acid,

- 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 15 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 20 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 25 3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-
35 2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

5 3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

10 Yet further illustrative are the compounds

3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15 3(S)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20

3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25 3(S)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
bis(trifluoroacetate) salt,

30

3(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

35 3(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

5 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

10

3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-
2-yl)propyl]imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

15

For use in medicine, the salts of the compounds of this invention refer to non-toxic "pharmaceutically acceptable salts." Other salts may, however, be useful in the preparation of the compounds according to the invention or of their pharmaceutically acceptable salts.

20 Salts encompassed within the term "pharmaceutically acceptable salts" refer to non-toxic salts of the compounds of this invention which are generally prepared by reacting the free base with a suitable organic or inorganic acid. Representative salts include the following: acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate,

25 bromide, calcium, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide, hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate,

30 mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, oxalate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, stearate, sulfate, subacetate, succinate, tannate, tartrate, teclate, tosylate, triethiodide and valerate.

35 Furthermore, where the compounds of the invention carry an acidic

moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, e.g., sodium or potassium salts; alkaline earth metal salts, e.g., calcium or magnesium salts; and salts formed with suitable organic ligands, e.g., quaternary ammonium salts.

5 The compounds of the present invention can have chiral centers and occur as racemates, racemic mixtures, diastereomeric mixtures, and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention. Therefore, where a compound is chiral, the separate enantiomers or
10 diastereomers, substantially free of the other, are included within the scope of the invention; further included are all mixtures of the two enantiomers. Also included within the scope of the invention are polymorphs and hydrates of the compounds of the instant invention.

 The present invention includes within its scope prodrugs of
15 the compounds of this invention. In general, such prodrugs will be functional derivatives of the compounds of this invention which are readily convertible *in vivo* into the required compound. Thus, in the methods of treatment of the present invention, the term "administering" shall encompass the treatment of the various conditions described with
20 the compound specifically disclosed or with a compound which may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to the patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in "Design of Prodrugs," ed. H. Bundgaard, Elsevier, 1985,
25 which is incorporated by reference herein in its entirety. Metabolites of these compounds include active species produced upon introduction of compounds of this invention into the biological milieu.

 The term "therapeutically effective amount" shall mean that amount of a drug or pharmaceutical agent that will elicit the
30 biological or medical response of a tissue, system, animal or human that is being sought by a researcher or clinician.

 The term "integrin receptor antagonist," as used herein, refers to a compound which binds to and antagonizes either the $\alpha\beta 3$ receptor, the $\alpha\beta 5$ receptor, or the $\alpha\beta 6$ receptor, or a compound which

binds to and antagonizes combinations of these receptors (for example, a dual $\alpha\text{v}\beta 3/\alpha\text{v}\beta 5$ receptor antagonist).

The term "bone resorption," as used herein, refers to the process by which osteoclasts degrade bone.

5 The term "alkyl" shall mean straight or branched chain alkanes of one to ten total carbon atoms, or any number within this range (i.e., methyl, ethyl, 1-propyl, 2-propyl, n-butyl, s-butyl, t-butyl, etc.).

10 The term "alkenyl" shall mean straight or branched chain alkenes of two to ten total carbon atoms, or any number within this range.

 The term "alkynyl" shall mean straight or branched chain alkynes of two to ten total carbon atoms, or any number within this range.

15 The term "cycloalkyl" shall mean cyclic rings of alkanes of three to eight total carbon atoms, or any number within this range (i.e., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl).

 The term "cycloheteroalkyl," as used herein, shall mean a 3- to 8-membered fully saturated heterocyclic ring containing one or two
20 heteroatoms chosen from N, O or S. Examples of cycloheteroalkyl groups include, but are not limited to piperidinyl, pyrrolidinyl, azetidiny, morpholinyl, piperazinyl.

 The term "alkoxy," as used herein, refers to straight or branched chain alkoxides of the number of carbon atoms specified (e.g.,
25 C₁₋₅ alkoxy), or any number within this range (i.e., methoxy, ethoxy, etc.).

 The term "aryl," as used herein, refers to a monocyclic or polycyclic system comprising at least one aromatic ring, wherein the monocyclic or polycyclic system contains 0, 1, 2, 3, or 4 heteroatoms
30 chosen from N, O, or S, and wherein the monocyclic or polycyclic system is either unsubstituted or substituted with one or more groups independently selected from hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₈ cycloalkyl, aryl, aryl C₁₋₈ alkyl, amino, amino C₁₋₈ alkyl, C₁₋₃ acylamino, C₁₋₃ acylamino C₁₋₈ alkyl, C₁₋₆ alkylamino, C₁₋₆ alkylamino
35 C₁₋₈ alkyl, C₁₋₆ dialkylamino, C₁₋₆ dialkylamino-C₁₋₈ alkyl, C₁₋₄ alkoxy,

- C1-4 alkoxy C1-6 alkyl, hydroxycarbonyl, hydroxycarbonyl C1-6 alkyl, C1-5 alkoxy carbonyl, C1-3 alkoxy carbonyl C1-6 alkyl, hydroxycarbonyl C1-6 alkyloxy, hydroxy, hydroxy C1-6 alkyl, cyano, trifluoromethyl, oxo or C1-5 alkylcarbonyloxy. Examples of aryl include, but are not limited to,
- 5 phenyl, naphthyl, pyridyl, pyrazinyl, pyrimidinyl, imidazolyl, benzimidazolyl, benzthiazolyl, benzoxazolyl, indolyl, thienyl, furyl, pyrrolyl, pyrazolyl, dihydrobenzofuryl, benzo(1,3) dioxolane, oxazolyl, isoxazolyl and thiazolyl, which are either unsubstituted or substituted with one or more groups independently selected from hydrogen,
- 10 halogen, C1-10 alkyl, C3-8 cycloalkyl, aryl, aryl C1-8 alkyl, amino, amino C1-8 alkyl, C1-3 acylamino, C1-3 acylamino C1-8 alkyl, C1-6 alkylamino, C1-6 alkylamino-C1-8 alkyl, C1-6 dialkylamino, C1-6 dialkylamino C1-8 alkyl, C1-4 alkoxy, C1-4 alkoxy C1-6 alkyl, hydroxycarbonyl, hydroxycarbonyl C1-6 alkyl, C1-5 alkoxy carbonyl, C1-3 alkoxy carbonyl
- 15 C1-6 alkyl, hydroxycarbonyl C1-6 alkyloxy, hydroxy, hydroxy C1-6 alkyl, cyano, trifluoromethyl, oxo or C1-5 alkylcarbonyloxy. Preferably, the aryl group is unsubstituted, mono-, di-, tri- or tetra-substituted with one to four of the above-named substituents; more preferably, the aryl group is unsubstituted, mono-, di- or tri-substituted with one to three of the
- 20 above-named substituents; most preferably, the aryl group is unsubstituted, mono- or di-substituted with one to two of the above-named substituents.

- Whenever the term "alkyl" or "aryl" or either of their prefix roots appears in a name of a substituent (e.g., aryl C0-8 alkyl), it shall be
- 25 interpreted as including those limitations given above for "alkyl" and "aryl." Designated numbers of carbon atoms (e.g., C1-10) shall refer independently to the number of carbon atoms in an alkyl or cyclic alkyl moiety or to the alkyl portion of a larger substituent in which alkyl appears as its prefix root.

- 30 The terms "arylalkyl" and "alkylaryl" include an alkyl portion where alkyl is as defined above and to include an aryl portion where aryl is as defined above. Examples of arylalkyl include, but are not limited to, benzyl, fluorobenzyl, chlorobenzyl, phenylethyl, phenylpropyl, fluorophenylethyl, chlorophenylethyl, thienylmethyl,

thienylethyl, and thienylpropyl. Examples of alkylaryl include, but are not limited to, toluene, ethylbenzene, propylbenzene, methylpyridine, ethylpyridine, propylpyridine and butylpyridine.

In the compounds of the present invention, two R¹ substituents, when on the same carbon atom, can be taken together with the carbon to which they are attached to form a carbonyl group.

In the compounds of the present invention, two R³ substituents, when on the same carbon atom, can be taken together with the carbon atom to which they are attached to form a carbonyl group. In such instances, the limitation, that in the resultant compound the carbon atom or atoms to which R³ is attached is itself attached to no more than one heteroatom, does not apply. Also, in the compounds of the present invention, two R³ substituents, when on the same carbon atom, can be taken together with the carbon atom to which they are attached to form a cyclopropyl group.

In the compounds of the present invention, R⁵ and R⁶ can be taken together with the carbon atom to which they are attached to form a carbonyl group. In such instances, the limitation, that in the resultant compound the carbon atom at which R⁵ and R⁶ is attached is itself attached to no more than one heteroatom, does not apply.

When substituents R⁷ and R⁸ include the definition C₀ (e.g., C₀₋₈ alkyl), the group modified by C₀ is not present in the substituent when C is zero. Similarly, when any of the variables m, n, t, or v, is zero, then the group modified by the variable is not present; for example, when t is zero, the group $-(CH_2)_tC\equiv CH$ is $-C\equiv CH$. In addition, the substituent "(C₁₋₆ alkyl)_pamino" where p is zero, one or two, refers to an amino, C₁₋₆ alkylamino and C₁₋₆ dialkylamino group, respectively. When a C₁₋₆ dialkylamino substituent is intended, the C₁₋₆ alkyl groups can be the same (e.g., dimethylamino) or different (e.g., N(CH₃)(CH₂CH₃)). Similarly, the substituent "(aryl)_pamino" or ["(aryl C₁₋₆ alkyl)_pamino"], where p is zero, one or two, refers to an amino, arylamino and diarylamino group, [or an amino, aryl C₁₋₆ alkylamino or di-(aryl C₁₋₆ alkyl)amino] respectively, where the aryl [or aryl C₁₋₆ alkyl] groups in a diarylamino [or di-(aryl C₁₋₆ alkyl)amino] substituent can be the same or different.


In the compounds of the present invention, R¹⁰ and R¹² can be taken together with the carbon atoms to which they are attached to form a 5- to 7-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of N, O, and S wherein said 5- to 7-membered monocyclic aromatic or nonaromatic ring system is either unsubstituted or substituted with one or more R¹ substituents.

The term "halogen" shall include iodine, bromine, chlorine, and fluorine.

The term "oxy" means an oxygen (O) atom. The term "thio" means a sulfur (S) atom. The term "oxo" means "=O." The term "carbonyl" means "C=O."

The term "substituted" shall be deemed to include multiple degrees of substitution by a named substituent. Where multiple substituent moieties are disclosed or claimed, the substituted compound can be independently substituted by one or more of the disclosed or claimed substituent moieties, singly or plurally. By independently substituted, it is meant that the (two or more) substituents can be the same or different.

Under standard nomenclature used throughout this disclosure, the terminal portion of the designated side chain is described first, followed by the adjacent functionality toward the point of attachment. For example, a C₁₋₅ alkylcarbonylamino C₁₋₆ alkyl substituent is equivalent to



In choosing compounds of the present invention, one of ordinary skill in the art will recognize that the various substituents, i.e. X, Y, Z, R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸, R⁹, R¹⁰, R¹¹, R¹², and R¹³ and the subscripts m, n, p, r, s, t, and v are to be chosen in conformity with well-known principles of chemical structure connectivity.

Representative compounds of the present invention typically display submicromolar affinity for the integrin receptors, particularly

the $\alpha v\beta 3$, $\alpha v\beta 5$, and/or $\alpha v\beta 6$ receptors. Compounds of this invention are therefore useful for treating mammals suffering from a bone condition caused or mediated by increased bone resorption, who are in need of such therapy. Pharmacologically effective amounts of the compounds, including pharmaceutically acceptable salts thereof, are administered to the mammal, to inhibit the activity of mammalian osteoclasts.

The compounds of the present invention are administered in dosages effective to antagonize the $\alpha v\beta 3$ receptor where such treatment is needed, as, for example, in the prevention or treatment of osteoporosis.

Further exemplifying the invention is the method wherein the integrin receptor antagonizing effect is an $\alpha v\beta 3$ antagonizing effect. An illustration of the invention is the method wherein the $\alpha v\beta 3$ antagonizing effect is selected from inhibition of bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, tumor growth, or metastasis. Preferably, the $\alpha v\beta 3$ antagonizing effect is the inhibition of bone resorption.

An example of the invention is the method wherein the integrin receptor antagonizing effect is an $\alpha v\beta 5$ antagonizing effect. More specifically, the $\alpha v\beta 5$ antagonizing effect is selected from inhibition of restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, tumor growth, or metastasis.

Illustrating the invention is the method wherein the integrin receptor antagonizing effect is a dual $\alpha v\beta 3/\alpha v\beta 5$ antagonizing effect. More particularly, the dual $\alpha v\beta 3/\alpha v\beta 5$ antagonizing effect is selected from inhibition of bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, tumor growth, or metastasis.

Illustrating the invention is the method wherein the integrin receptor antagonizing effect is an $\alpha v\beta 6$ antagonizing effect. More particularly, the $\alpha v\beta 6$ antagonizing effect is selected from inhibition of angiogenesis, inflammatory response, or wound healing.

Illustrating the invention is the method wherein the $\alpha v\beta 3$ antagonizing effect is selected from inhibition of bone resorption,

inhibition of restenosis, inhibition of angiogenesis, inhibition of diabetic retinopathy, inhibition of macular degeneration, inhibition of atherosclerosis, inflammation, viral disease, or inhibition of tumor growth or metastasis. Preferably, the $\alpha v \beta 3$ antagonizing effect is the inhibition of bone resorption.

More particularly illustrating the invention is a pharmaceutical composition comprising any of the compounds described above and a pharmaceutically acceptable carrier. Another example of the invention is a pharmaceutical composition made by combining any of the compounds described above and a pharmaceutically acceptable carrier. Another illustration of the invention is a process for making a pharmaceutical composition comprising combining any of the compounds described above and a pharmaceutically acceptable carrier.

Further illustrating the invention is a method of treating and/or preventing a condition mediated by antagonism of an integrin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds described above. Preferably, the condition is selected from bone resorption, osteoporosis, restenosis, diabetic retinopathy, macular degeneration, angiogenesis, atherosclerosis, inflammation, viral disease, cancer, tumor growth, and metastasis. More preferably, the condition is selected from osteoporosis and cancer. Most preferably, the condition is osteoporosis.

More specifically exemplifying the invention is a method of eliciting an integrin antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above. Preferably, the integrin antagonizing effect is an $\alpha v \beta 3$ antagonizing effect; more specifically, the $\alpha v \beta 3$ antagonizing effect is selected from inhibition of bone resorption, inhibition of restenosis, inhibition of atherosclerosis, inhibition of angiogenesis, inhibition of diabetic retinopathy, inhibition of macular degeneration, inhibition of inflammation, inhibition of viral disease, or inhibition of tumor growth or metastasis. Most preferably, the $\alpha v \beta 3$

antagonizing effect is inhibition of bone resorption. Alternatively, the integrin antagonizing effect is an $\alpha v \beta 5$ antagonizing effect, an $\alpha v \beta 6$ antagonizing effect, or a mixed $\alpha v \beta 3$, $\alpha v \beta 5$, and $\alpha v \beta 6$ antagonizing effect. Examples of $\alpha v \beta 5$ antagonizing effects are inhibition of

- 5 restenosis, atherosclerosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, or tumor growth. Examples of dual $\alpha v \beta 6$ antagonizing effects are inhibition of angiogenesis, inflammatory response and wound healing.

- Additional examples of the invention are methods of
10 inhibiting bone resorption and of treating and/or preventing osteoporosis in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of any of the compounds or any of the pharmaceutical compositions described above.

- Additional illustrations of the invention are methods of
15 treating hypercalcemia of malignancy, osteopenia due to bone metastases, periodontal disease, hyperparathyroidism, periarticular erosions in rheumatoid arthritis, Paget's disease, immobilization-induced osteopenia, and glucocorticoid treatment in a mammal in need thereof, comprising administering to the mammal a therapeutically
20 effective amount of any of the compounds or any of the pharmaceutical compositions described above.

- More particularly exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment and/or prevention of osteoporosis in a
25 mammal in need thereof. Still further exemplifying the invention is the use of any of the compounds described above in the preparation of a medicament for the treatment and/or prevention of bone resorption, tumor growth, cancer, restenosis, atherosclerosis, diabetic retinopathy, macular degeneration, inflammation, viral disease, and/or
30 angiogenesis.

Also exemplifying the invention are compositions further comprising an active ingredient selected from the group consisting of

- a.) an organic bisphosphonate or a pharmaceutically acceptable salt or ester thereof,
35 b.) an estrogen receptor modulator,

- c.) a cytotoxic/antiproliferative agent,
- d.) a matrix metalloproteinase inhibitor,
- e.) an inhibitor of epidermal-derived, fibroblast-derived, or platelet-derived growth factors,
- 5 f.) an inhibitor of VEGF,
- g.) an inhibitor of Flk-1/KDR, Flt-1, Tck/Tie-2, or Tie-1,
- h.) a cathepsin K inhibitor; and
- i.) a prenylation inhibitor, such as a farnesyl transferase inhibitor or a geranylgeranyl transferase inhibitor or a dual
- 10 farnesyl/geranylgeranyl transferase inhibitor; and mixtures thereof.

(See, B. Millauer et al., "Dominant-Negative Inhibition of Flk-1 Suppresses the Growth of Many Tumor Types in *Vivo*", Cancer Research, 56, 1615-1620 (1996), which is incorporated by reference herein

15 in its entirety).

Preferably, the active ingredient is selected from the group consisting of:

- a.) an organic bisphosphonate or a pharmaceutically acceptable salt or ester thereof,
- 20 b.) an estrogen receptor modulator, and
- c.) a cathepsin K inhibitor; and mixtures thereof.

Nonlimiting examples of such bisphosphonates include alendronate, etidronate, pamidronate, risedronate, ibandronate, and pharmaceutically acceptable salts and esters thereof. A particularly

25 preferred bisphosphonate is alendronate, especially alendronate monosodium trihydrate.

Nonlimiting examples of estrogen receptor modulators include estrogen, progesterin, estradiol, droloxifene, raloxifene, and tamoxifene.

30 Nonlimiting examples of cytotoxic/antiproliferative agents are taxol, vincristine, vinblastine, and doxorubicin.

Cathepsin K, formerly known as cathepsin O2, is a cysteine protease and is described in PCT International Application Publication No. WO 96/13523, published May 9, 1996; U.S. Patent No. 5,501,969,

35 issued March 3, 1996; and U.S. Patent No. 5,736,357, issued April 7, 1998,

all of which are incorporated by reference herein in their entirety. Cysteine proteases, specifically cathepsins, are linked to a number of disease conditions, such as tumor metastasis, inflammation, arthritis, and bone remodeling. At acidic pH's, cathepsins can degrade type-I collagen. Cathepsin protease inhibitors can inhibit osteoclastic bone resorption by inhibiting the degradation of collagen fibers and are thus useful in the treatment of bone resorption diseases, such as osteoporosis.

The present invention is also directed to combinations of the compounds of the present invention with one or more agents useful in the prevention or treatment of osteoporosis. For example, the compounds of the instant invention may be effectively administered in combination with effective amounts of other agents such as an organic bisphosphonate, an estrogen receptor modulator, or a cathepsin K inhibitor.

Additional illustrations of the invention are methods of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound described above and one or more agents known to be cytotoxic/antiproliferative. Also, the compounds of the present invention can be administered in combination with radiation therapy for treating tumor growth and metastasis.

In addition, the integrin $\alpha v \beta 3$ antagonist compounds of the present invention may be effectively administered in combination with a growth hormone secretagogue in the therapeutic or prophylactic treatment of disorders in calcium or phosphate metabolism and associated diseases. These diseases include conditions which can benefit from a reduction in bone resorption. A reduction in bone resorption should improve the balance between resorption and formation, reduce bone loss or result in bone augmentation. A reduction in bone resorption can alleviate the pain associated with osteolytic lesions and reduce the incidence and/or growth of those lesions. These diseases include: osteoporosis (including estrogen deficiency, immobilization, glucocorticoid induced and senile), osteodystrophy, Paget's disease, myositis ossificans, Bechterew's disease, malignant hypercalcemia, metastatic bone disease, periodontal disease,

cholelithiasis, nephrolithiasis, urolithiasis, urinary calculus, hardening of the arteries (sclerosis), arthritis, bursitis, neuritis and tetany. Increased bone resorption can be accompanied by pathologically high calcium and phosphate concentrations in the plasma, which would be alleviated by this treatment. Similarly, the present invention would be useful in increasing bone mass in patients with growth hormone deficiency. Thus, preferred combinations are simultaneous or alternating treatments of an $\alpha\text{v}\beta 3$ receptor antagonist of the present invention and a growth hormone secretagogue, optionally including a third component comprising an organic bisphosphonate, preferably alendronate monosodium trihydrate.

In accordance with the method of the present invention, the individual components of the combination can be administered separately at different times during the course of therapy or concurrently in divided or single combination forms. The instant invention is therefore to be understood as embracing all such regimes of simultaneous or alternating treatment, and the term "administering" is to be interpreted accordingly. It will be understood that the scope of combinations of the compounds of this invention with other agents useful for treating integrin-mediated conditions includes in principle any combination with any pharmaceutical composition useful for treating osteoporosis.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts.

The compounds of the present invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions, syrups and emulsions. Likewise, they may also be administered in intravenous (bolus or infusion), intraperitoneal, topical (e.g., ocular eyedrop), subcutaneous, intramuscular or transdermal (e.g., patch) form, all using forms well known to those of ordinary skill in the pharmaceutical

arts. An effective but non-toxic amount of the compound desired can be employed as an $\alpha\text{v}\beta 3$ antagonist.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including
5 type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of administration; the renal and hepatic function of the patient; and the particular compound or salt thereof employed. An ordinarily skilled physician, veterinarian
10 or clinician can readily determine and prescribe the effective amount of the drug required to prevent, counter or arrest the progress of the condition.

Oral dosages of the present invention, when used for the indicated effects; will range between about 0.01 mg per kg of body weight per day (mg/kg/day) to about 100 mg/kg/day, preferably 0.01 to 10
15 mg/kg/day, and most preferably 0.1 to 5.0 mg/kg/day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. A medicament
20 typically contains from about 0.01 mg to about 500 mg of the active ingredient, preferably, from about 1 mg to about 100 mg of active ingredient. Intravenously, the most preferred doses will range from about 0.1 to about 10 mg/kg/minute during a constant rate infusion. Advantageously, compounds of the present invention may be
25 administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, preferred compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal
30 skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

In the methods of the present invention, the compounds
35 herein described in detail can form the active ingredient, and are

typically administered in admixture with suitable pharmaceutical diluents, excipients or carriers (collectively referred to herein as 'carrier' materials) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine or phosphatidylcholines.

Compounds of the present invention may also be delivered by the use of monoclonal antibodies as individual carriers to which the compound molecules are coupled. The compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol, polyhydroxyethylaspartamide-phenol, or polyethyleneoxide-polylysine substituted

with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, 5 polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihydropyrans, polycyanoacrylates and crosslinked or amphipathic block copolymers of hydrogels.

In the schemes and examples below, various reagent
 10 symbols and abbreviations have the following meanings:

AcOH:	Acetic acid.
BH ₃ •DMS:	Borane•dimethylsulfide.
BOC(Boc):	t-Butyloxycarbonyl.
BOP:	Benzotriazol-1-yloxytris(dimethylamino)-
15	phosphonium hexafluorophosphate.
CBZ(Cbz):	Carbobenzyloxy or benzyloxycarbonyl.
CDI:	Carbonyldiimidazole.
CH ₂ Cl ₂ :	Methylene chloride.
CH ₃ CN:	Acetonitrile
20	CHCl ₃ : Chloroform.
DBA:	Bis(dibenzylidene)acetone.
DEAD:	Diethyl azodicarboxylate.
DIAD:	Diisopropyl azodicarboxylate.
DIBAH or	
25	DIBAL-H: Diisobutylaluminum hydride.
DIPEA:	Diisopropylethylamine.
DMAP:	4-Dimethylaminopyridine.
DME:	1,2-Dimethoxyethane.
DMF:	Dimethylformamide.
30	DMSO: Dimethylsulfoxide.
DPPF:	1,1'-bis(diphenylphosphino)ferrocene.
DPFN:	3,5-Dimethyl-1-pyrazolylformamidinium nitrate.
EDC:	1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide •HCl
EtOAc:	Ethyl acetate.
35	EtOH: Ethanol.

	HOAc:	Acetic acid.
	HOAT:	1-Hydroxy-7-azabenzotriazole
	HOBT:	1-Hydroxybenzotriazole.
	IBCF:	Isobutylchloroformate
5	LDA:	Lithium diisopropylamide.
	MeOH:	Methanol.
	MMNG	1,1-methyl-3-nitro-1-nitrosoguanidine
	NEt ₃ :	Triethylamine.
	NMM:	N-methylmorpholine.
10	PCA • HCl:	Pyrazole carboxamidine hydrochloride.
	Pd/C:	Palladium on activated carbon catalyst.
	Ph:	Phenyl.
	pTSA	p-Toluenesulfonic acid.
	TEA:	Triethylamine.
15	TFA:	Trifluoroacetic acid.
	THF:	Tetrahydrofuran.
	TLC:	Thin Layer Chromatography.
	TMEDA:	N,N,N',N'-Tetramethylethylenediamine.
	TMS:	Trimethylsilyl.

20 The novel compounds of the present invention can be prepared according to the procedure of the following schemes and examples, using appropriate materials and are further exemplified by the following specific examples. The compounds illustrated in the examples are not, however, to be construed as forming the only genus
25 that is considered as the invention. The following examples further illustrate details for the preparation of the compounds of the present invention. Those skilled in the art will readily understand that known variations of the conditions and processes of the following preparative procedures can be used to prepare these compounds. All temperatures
30 are degrees Celsius unless otherwise noted.


 The following Schemes and Examples describe procedures for making representative compounds of the present invention. Moreover, by utilizing the procedures described in detail in PCT International Application Publication Nos. WO95/32710, published 7
35 December 1995, and WO95/17397, published 29 June 1995, both of which

are incorporated by reference herein in their entirety, in conjunction with the disclosure contained herein, one of ordinary skill in the art can readily prepare additional compounds of the present invention claimed herein. Additionally, for a general review describing the synthesis of

5 β -alanines which can be utilized as the C-terminus of the compounds of the present invention, see Cole, D.C., *Recent Stereoselective Synthetic Approaches to β -Amino Acids*, *Tetrahedron*, 1994, 50, 9517-9582; Juaristi, E, *et al.*, *Enantioselective Synthesis of β -Amino Acids*, *Aldrichimica Acta*, 1994, 27, 3. In particular, synthesis of the 3-methyl- β -alanine is

10 taught in Duggan, M.F. *et al.*, *J. Med. Chem.*, 1995, 38, 3332-3341; the 3-ethynyl- β -alanine is taught in Zablocki, J.A., *et al.*, *J. Med. Chem.*, 1995, 38, 2378-2394; the 3-(pyridin-3-yl)- β -alanine is taught in Rico, J.G. *et al.*, *J. Org. Chem.*, 1993, 58, 7948-7951; and the 2-amino- and 2-tosylamino- β -alanines are taught in Xue, C-B, *et al.*, *Biorg. Med. Chem. Letts.*, 1996,

15 6, 339-344. The references described in this paragraph are all also incorporated by reference herein in their entirety.

SCHEME 1

1-Bromo-3-(2,2-diethoxy-ethoxy)-benzene (1-2)

To a suspension of NaH (2.77 g, 115.6 mmol) in DMF (100 mL) at 0°C was added a solution of 3-bromophenol 1-1 in DMF (40 mL) over 40 min. After the addition was complete, the solution was stirred
5 for an additional 30 min. The solution was then treated with neat bromoacetaldehyde diethyl acetal (17.36 g, 115.6 mmol). The solution was heated at 100°C for 8 h, cooled to room temperature, and extracted with Et₂O (3 x 200 mL). The combined organic extracts were washed with 10% aq. NaOH (100 mL) and brine (100 mL), dried over MgSO₄,
10 filtered and concentrated to give 1-2 as a yellow oil.
TLC R_f = 0.4 (10% ethyl acetate/hexanes).
¹H NMR (300 MHz, CDCl₃) δ 7.19-7.05 (m, 3H), 6.85 (d, 1H), 4.81 (t, 1H, J=6.8 Hz), 3.99 (d, 2H, J=6.8 Hz), 3.71 (m, 4H), 1.22 (t, 6H, J=7.1 Hz)

6-Bromo-benzofuran (1-3)

To a solution of the acetal 1-2 in toluene (200 mL) was added polyphosphoric acid (20 g). The biphasic mixture was heated to 100°C and stirred at this temperature for 4 h. The mixture was cooled to room temperature, poured onto ice, and extracted with Et₂O (2 x 200 mL). The
20 combined organic extracts were washed with saturated aq. NaHCO₃ and brine. The solution was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (100% hexanes) to give the product 1-3 as a yellow oil.
TLC R_f = 0.3 (100% hexanes).
25 ¹H NMR (300 MHz, CDCl₃) δ 7.68 (s, 1H), 7.60 (d, 1H, J=2.1 Hz), 7.46 (d, 1H, J=8.4 Hz), 7.36 (dd, 1H, J=8.1, 1.5 Hz), 6.75 (dd, 1H, J=7.1, 0.9 Hz).

3-Benzofuran-6-yl-acrylic acid ethyl ester (1-4)

A mixture of the 6-bromo-benzofuran 1-3 (1.74 g, 8.79
30 mmol), ethyl acrylate (1.09 g, 10.98 mmol), Pd(OAc)₂ (0.099 g, 0.44 mmol), tri-*o*-tolylphosphine (0.268 g, 0.880 mmol), and sodium acetate (3.60 g, 43.9 mmol) in DMF (10 mL) was heated to 100°C in a sealed tube for 4 h. The mixture was cooled to room temperature, diluted with water, and extracted with Et₂O (2 x 40 mL). The combined organic
35 extracts were washed with brine (30 mL), dried over MgSO₄, filtered,

and concentrated. The residue was purified by flash chromatography (10% ethyl acetate/hexanes) to give the ester 1-4 as an off-white solid.

TLC Rf = 0.3 (10% ethyl acetate/hexanes).

¹H NMR (300 MHz, CDCl₃) δ 7.78 (d, 1H, J=15.9 Hz), 7.68 (d, 1H, J=2.4 Hz), 7.66 (s, 1H), 7.59 (d, 1H, J=8.4 Hz), 7.43 (dd, 1H, J=9.0, 1.5 Hz), 6.78 (m, 1H), 6.47 (d, 1H, J=15.9 Hz), 4.27 (q, 2H, J=7.2 Hz), 1.34 (t, 3H, J=7.2 Hz).

10 3(S)-Benzofuran-6-yl-3-[benzyl-(1(R)-phenethyl)-amino]-propionic acid ethyl ester (1-5)

A solution of N-benzyl-α-(R)-methylbenzylamine (1.32 g, 6.30 mmol) in THF (25 mL) at 0°C was treated with n-BuLi (2.52 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0°C for 30 min and then cooled to -78°C. A solution of acrylate 1-4 (0.681 g, 3.15 mmol) in THF (5 mL) was added. After stirring for 15 min at -78°C, satd. aq. NH₄Cl soln (5 mL) was added and the cold bath removed. The mixture was warmed to room temperature and extracted with Et₂O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL), dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (10% ethyl acetate/hexanes) to give the β-aminoester 1-5 as a yellow oil.

TLC Rf = 0.8 (10% ethanol/dichloromethane).

¹H NMR (300 MHz, CDCl₃) δ 7.58 (m, 3H), 7.41 (m, 2H), 7.22 (m, 9H), 7.59 (s, 1H), 4.58 (m, 1H), 4.05 (m, 1H), 3.91 (q, 2H, J=7.1 Hz), 3.72 (m, 2H), 2.62 (m, 2H), 1.21 (d, 3H, J=7.2 Hz), 1.03 (t, 3H, J=7.1 Hz).

3(S)-Amino-3-(2,3-dihydro-benzofuran-6-yl)-propionic acid ethyl ester (1-6)


A mixture of the dibenzylamine 1-5 (1.19 g, 2.78 mmol) in EtOH/H₂O/AcOH (26 mL/3 mL/1.0 mL) was degassed with argon and treated with Pd(OH)₂ (1.19 g). The mixture was placed under 1 atm of H₂. After stirring for 18 h, the mixture was diluted with EtOAc, and filtered through celite. The filtrate was concentrated and the residue purified by flash chromatography (10% ethyl acetate/dichloromethane) to give the ester 1-6 as a white solid.


TLC Rf = 0.25 (10% ethanol/dichloromethane).


^1H NMR (300 MHz, CD_3OD) as the trifluoroacetate salt: δ 7.25 (d, 1H, $J=8.1$ Hz), 6.88 (m, 1H), 7.66 (s, 1H), 6.82 (s, 1H), 4.58 (m, 3H), 4.12 (m, 2H), 3.30 (m, 1H), 3.19 (m, 2H), 2.98 (m, 2H), 1.11 (t, 3H, $J=7.2$ Hz).

5

SCHEME 2







4-Oxo-pentanoic acid methoxy-methyl-amide (2-1)

To a stirred solution of levulinic acid (30 g, 0.258 mol) in
 5 CHCl_3 (850 mL) at 0°C was added triethylamine (43.2 mL, 0.310 mol),
 followed by isobutyl chloroformate (37 mL, 0.284 mol) over 15 minutes.
 After 30 minutes, triethylamine (57.6 mL, 0.413 mol) was added, followed
 by N,O-dimethylhydroxylamine hydrochloride (37.8 g, 0.387 mol) in 5
 portions over 5 minutes. Vigorous bubbling ensued, and the mixture
 10 was allowed to warm to RT and stirred for 1 h. The mixture was
 reduced to a moist solid by rotary evaporation under reduced pressure,

slurried in 500 mL EtOAc, washed with 10% K₂CO₃, brine, and dried over Na₂SO₄. Evaporative removal of the solvent gave 2-1 as a yellow oil. TLC R_f = 0.42 (silica, 1:1 chloroform/ethyl acetate).

¹H NMR (300 MHz, CDCl₃) δ 3.74 (s, 3H), 3.18 (s, 3H), 2.65-2.95 (m, 4H),
5 2.21 (s, 3H).

N-methoxy-N-methyl-3-(2-methyl-[1,3]dioxolan-2-yl)-propionamide (2-2)

To a solution of 2-1 (38 g, 0.239 mol) in 500 mL benzene was added ethylene glycol (17.3 mL, 0.310 mol) and p-toluenesulfonic acid (1
10 g). The mixture was heated at reflux for 2 h with azeotropic removal of water. After cooling, the solution was washed with 200 mL sat. NaHCO₃, brine, and dried over Na₂SO₄. Evaporative removal of the solvent gave 2-2 as a yellow oil.

TLC R_f = 0.62 (silica, ethyl acetate).

15 ¹H NMR (300 MHz, CDCl₃) δ 3.95 (m, 4H), 3.68 (s, 3H), 3.17 (s, 3H), 2.51 (t, 2H, J=8 Hz), 2.00 (t, 3H, J=6 Hz) 1.33 (s, 3H).

3-(2-Methyl-[1,3]dioxolan-2-yl)-propionaldehyde (2-3)

To a solution of 2-2 (44.74 g, 0.22 mol) in 400 mL THF at
20 -78°C was added DIBAL (264 mL 1 M in hexanes, 0.264 mol) over 10 minutes. After stirring for 1 h, 350 mL of 1.0 M Rochelle's salt and 300 mL ether were added followed by the removal of the cooling bath. After stirring for 1 h, the organic portion was separated and dried over Na₂SO₄. Evaporative removal of the solvent gave 2-3 as a colorless oil.

25 TLC R_f = 0.80 (silica, ethyl acetate).

¹H NMR (300 MHz, CDCl₃) δ 9.73 (s, 1H), 3.50 (d, 1H, J=16 Hz), 2.61 (d, 1H, J=21 Hz), 2.48 (m, 1H), 2.07 (t, 1H, J=7H), 1.33 (s, 3H).

[3-(2-Methyl-[1,3]dioxolan-2-yl)-propylamino]-acetic acid ethyl ester (2-4)

30 To a solution of 2-3 (31.7 g, 0.22 mol) in 1000 mL 1,2-dichloroethane at 0°C were added glycine ethyl ester hydrochloride (61.5 g, 0.44 mol), triethylamine (107 mL, 0.77 mol), and NaB(OAc)₃H (65.3 g, 0.308 mol). The mixture was allowed to warm to RT and stirred for 15 h. The mixture was evaporated to one-third its initial volume, diluted with

EtOAc and then washed with 10% K₂CO₃, brine, and dried over Na₂SO₄. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 1:1 chloroform/ethyl acetate followed by 5% MeOH/ethyl acetate) to give 2-4 as a yellow oil.

5 TLC R_f = 0.40 (silica, ethyl acetate).

¹H NMR (300 MHz, CDCl₃) δ 6.01 (br s, 1H), 4.21 (m, 3H), 4.03 (d, 1H, J=5 Hz), 3.93 (m, 4H), 2.62 (t, 2H, J=8 Hz), 1.53-1.67 (m, 4H), 1.29 (m, 6H).

10 [Tert-butoxycarbonyl-[3-(2-methyl-[1,3]dioxolan-2-yl)-propyl]-amino]-acetic acid ethyl ester (2-5)

To a solution of 2-4 (24 g, 0.104 mol) in 100 mL THF were added a trace of DMAP, 20 drops of triethylamine, and BOC₂O (23.8 g, 0.109 mol). After 4 h, evaporative removal of the solvent gave 2-5 as a colorless oil.

15 TLC R_f = 0.38 (silica, 30% ethyl acetate/hexane).

¹H NMR (300 MHz, CDCl₃, mixture of rotamers) δ 4.22 (m, 3H), 3.93 (m, 4H), 3.27 (m, 2H), 1.63 (m, 4H), 1.51 (s, 3H), 1.47 (s, 3H), 1.42 (s, 3H), 1.31 (s, 3H), 1.28 (m, 4H).

20 [Tert-butoxycarbonyl-(4-oxo-pentyl)-amino]-acetic acid ethyl ester (2-6)

To a solution of 2-5 (35 g, 0.1 mol) in 600 mL acetone was added p-toluenesulfonic acid (1 g). The mixture was heated at reflux for 2 h. After cooling, the mixture was evaporated to one-fifth its initial volume, diluted with EtOAc and then washed with sat. NaHCO₃, brine, and dried over Na₂SO₄. Evaporative removal of the solvent gave 2-6 as a yellow oil.

25 TLC R_f = 0.31 (silica, 30% ethyl acetate/hexane).

¹H NMR (300 MHz, CDCl₃, mixture of rotamers) δ 4.20 (m, 2H), 3.92 (s, 0.85H), 3.83 (s, 1.15 H), 3.3 (m, 2H), 2.52 (m, 2H), 2.14 (s, 3H), 1.78 (m, 2H), 1.51-1.42 (3s, 9H), 1.28 (m, 3H).

30

[Tert-butoxycarbonyl-(3-[1,8]naphthyridin-2-yl-propyl)-amino]-acetic acid ethyl ester (2-7)

A solution of 2-6 (28 g, 97.4 mmol), 2-amino-3 formylpyridine (15.5 g, 127 mmol), proline (11.2 g, 97.4 mmol) in ethanol (250 mL) was heated at reflux for 15 h. After cooling and evaporation, the residue was chromatographed (silica gel, 1:1 chloroform/ethyl acetate) to give 2-7 as a yellow oil.

TLC R_f = 0.41 (silica, 70:25:5 chloroform/ethyl acetate/methanol)

¹H NMR (300 MHz, CDCl₃) δ 9.09 (m, 1H), 8.14 (m, 2H), 7.43 (m, 2H), 4.17 (q, 2H, 7 Hz), 3.9 (2s, 2H), 3.43 (q, 2H, J=7 Hz), 3.07 (m, 2H), 2.18 (m, 2H), 1.42 (s, 9H), 1.25 (m, 3H).

[(Tert-butoxycarbonyl-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino)-acetic acid ethyl ester (2-8)]

A solution of 2-7 (24.3 g, 65.1 mmol), platinum oxide (4 g) and ethanol (130 mL) was stirred under a balloon of hydrogen gas for 6 h. Following filtration and evaporation, the residue was chromatographed (silica gel, ethyl acetate) to give 2-8 as a yellow oil.

TLC R_f = 0.35 (silica, 70:25:5 chloroform/ethyl acetate/methanol)

¹H NMR (300 MHz, CDCl₃) δ 7.05 (d, 1H, J=6 Hz), 6.37 (m, 1H), 4.74 (brs, 1H), 4.18 (q, 2H, J=7 Hz), 3.9 (2s, 2H), 3.32 (m, 4H), 2.63 (m, 2H), 2.51 (m, 2H), 2.72 (m, 4H), 1.43 (m, 9H), 1.26 (m, 3H).

[(Methoxy-methyl-carbamoyl)-methyl]-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-carbamic acid tert-butyl ester (2-9)]

To a solution of 2-8 (1.49 g, 3.96 mmol) in ethanol (8 mL) was added NaOH (4.36 mL 1M solution in water, 4.36 mmol). After stirring for 1 h at 50°C, HCl (4.75 mL of a 1M solution in water, 4.75 mmol) was added, and the mixture evaporated to give an oily residue. The residue was evaporated from ethanol three times, and then from acetonitrile three times, producing a yellow crusty solid which was dried under a vacuum of <2 mm Hg for 2 h. This residue was then slurried in chloroform (15 mL), and triethylamine (2.75 mL, 19.8 mmol), N,O-dimethylhydroxylamine hydrochloride (0.772 g, 7.92 mmol), HOBT (1g) and EDC (0.91 g, 4.75 mmol) were added. After stirring for 15 h, the mixture was evaporated to dryness, the residue slurried in EtOAc,

washed with sat. NaHCO₃, brine, and dried over Na₂SO₄. Evaporative removal of the solvent gave 2-9 as a yellow oil.

TLC R_f = 0.49 (silica, 70:25:5 chloroform /ethyl acetate /methanol)

¹H NMR (300 MHz, CDCl₃) δ 7.05 (m, 1H), 6.38 (m, 1H), 4.81 (br s, 1H),
5 3.69, m, 3H), 3.37 (m, 4H), 3.18 (s, 3H), 2.64 (m, 2H), 2.53 (m, 2H), 1.88 (m, 4H), 1.44 (m, 9H).

{Tert-butoxycarbonyl-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino}-acetaldehyde (2-9A)

10 To a stirred solution of 2-9 (11.0 g, 28.0 mmol) and THF (300 ml) at -78°C was added DIBAL (1.0M/hexanes, 42 ml, 42 mmol) dropwise over 20 minutes. After 1.0 hour, 300 ml of 1.0 M Rochelle's salt was added followed by the removal of the cooling bath. The mixture was stirred for 1.0 hour and then diluted with Et₂O. After 30 minutes of
15 stirring, the organic portion was separated and dried over MgSO₄.

Evaporative removal of the solvent gave crude aldehyde 2-9A as a colorless oil.

TLC R_f = 0.34 (silica, 75:25:5 chloroform /EtOAc/MeOH).

20 3(S)-(2-(Tert-butoxycarbonyl-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino)-ethylamino)-3-(2,3-dihydro-benzofuran-6-yl)-propionic acid ethyl ester (2-10)

A mixture of the crude aldehyde 2-9A (9.1 mmol), 1-6 (3.2 g, 11.8 mmol), powdered molecular sieves (3 g) and DCE (100 mL) was
25 stirred for 30 minutes. The mixture was cooled to 0°C and then NaB(OAc)₃H (2.7 g, 12.7 mmol) was added. After 1 hour, the reaction was diluted with EtOAc and then washed with 10% K₂CO₃, brine, and dried over MgSO₄. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 1-3% [10:10:1 EtOH/ NH₄OH/
30 H₂O]/50:50 chloroform /ethyl acetate) to give 2-10 as a yellow oil.

TLC R_f = 0.23 (silica, 5% [10:10:1 EtOH/ NH₄OH/ H₂O]/50:50 chloroform /ethyl acetate)

¹H NMR (300 MHz, CDCl₃) δ 7.10 (d, J=7.6 Hz, 1H), 7.04 (d, J=7.3 Hz, 1H),
6.75 (m, 2H), 6.31 (d, J=7.3 Hz, 1H), 4.76 (s, 1H), 4.55 (t, J=8.6 Hz, 2H), 4.08
35 (m, 2H), 4.00 (t, J=6.1 Hz, 1H), 3.41 (m, 2H), 3.16 (m, 6H), 2.68 (t, J=6.4 Hz,

1H), 2.59 (m, 3H), 2.48 (t, J=7.6 Hz, 2H), 1.81 (m, 4H), 1.39 (s, 9H), 1.21 (m, 3H).

5 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propylamino]-ethylamino}-propionic acid ethyl ester (2-11)

HCl gas was rapidly bubbled through a solution of 2-10 (4.0 g, 7.2 mmol) in dioxane (160 ml) at 0°C for 10 minutes. After 30 minutes, the solution was purged with argon for 30 minutes. The solution was concentrated to give the amine 2-11 as a yellow solid.

10 ¹H NMR (300 MHz, CDCl₃) δ 7.91 (s, 1H), 7.40 (d, J=7.0 Hz, 1H), 7.23 (d, J=7.6 Hz, 1H), 7.10 (m, 2H), 6.56 (d, J=6.1 Hz, 1H), 4.58 (m, 2H), 4.04 (m, 2H), 3.49 (m, 4H), 3.19 (m, 4H), 2.90 (m, 2H), 2.79 (m, 2H), 2.30 (m, 2H), 1.98 (m, 2H), 1.85 (m, 5H), 1.15 (t, J=7.1 Hz, 3H).

15 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl ester (2-12)

To a stirred mixture of 2-11 (11.8 mmol), CH₂Cl₂ (3 mL) and 20 20% K₂CO₃ was added phosgene (1.93 M toluene, 6.7 ml, 13.0 mmol) dropwise over 20 minutes. After stirring for 30 minutes, the organic layer was separated and dried over MgSO₄. Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 5-10% methanol /ethyl acetate) to give 2-12 as a yellow oil.

25 TLC R_f = 0.25 (silica, 70:20:10 chloroform /ethyl acetate /methanol)
¹H NMR (300 MHz, CDCl₃) δ 7.12 (d, J=7.6 Hz, 1H), 7.02 (d, J=7.3 Hz, 1H), 6.80 (d, J=7.6 Hz, 1H), 6.70 (s, 1H), 6.34 (d, J=7.3 Hz, 1H), 5.46 (t, J=7.9 Hz, 1H), 4.74 (s, 1H), 4.55 (t, J=8.9 Hz, 2H), 4.10 (q, J=7.3 Hz, 2H), 3.41 (m, 2H), 3.21 (m, 6H), 2.95 (m, 3H), 2.67 (t, J=6.1 Hz, 2H), 2.52 (t, J=7.6 Hz, 30 2H), 1.88 (m, 5H), 1.20 (t, J=7.3 Hz, 3H).




3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (2-13)


To a solution of 2-12 (2.9g, 6.06 mmol) in EtOH (15 mL) was 35 added 1N NaOH (7.2 ml, 7.2 mmol). After stirring for 2 h, the solvents

were evaporated and the residue chromatographed (silica gel, 25:10:1:1 followed by 15:10:1:1 ethyl acetate /EtOH /water /NH₄OH) to give 2-13 as a white solid.

TLC R_f = 0.24 (15:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

- 5 ¹H NMR (300 MHz, CD₃OD) δ 7.55 (d, J=7.3 Hz, 1H), 7.18 (d, J=7.6 Hz, 1H), 6.81 (d, J=6.1 Hz, 1H), 6.72 (s, 1H), 6.62 (d, J=7.3 Hz, 1H), 5.38 (t, J=7.9 Hz, 1H), 4.53 (t, J=8.9 Hz, 2H), 3.14 → 3.53 (9H), 2.97 (m, 3H), 2.80 (t, J=6.1 Hz, 2H), 2.67 (t, J=7.3 Hz, 2H), 1.93 (m, 4H).

SCHEME 3

SCHEME 3 (CONTINUED)

5

Ethyl 3-fluorocinnamate (3-2)

To a solution of 3-fluorobenzaldehyde 3-1 (18.16 g, 146 mmol) in dichloromethane (500 mL) was added ethyl
 10 (triphenylphosphoranylidene)acetate (61.2 g, 176 mmol) and the resulting solution was stirred at room temperature for 18 hr. After evaporation of the solvent, the residue was swirled with ether/hexane and filtered. The filtrate was concentrated and then purified on a plug of silica gel eluting with hexane/EtOAc 9:1. Removal of the solvent
 15 afforded the title compound 3-2 as an oil (~95% trans) which was used without further purification in the next step.

^1H NMR (300 MHz, CDCl_3) δ 1.36 (3H, t), 4.28 (2H, q), 6.43 (1H, d), 7.08 (1H, m), 7.2-7.4 (3H, m), 7.64 (1H, d).

20 N-Benzyl-(R)- α -methylbenzyl-3(S)-fluorophenyl- β -alanine ethyl ester (3-3)

To a solution of N-benzyl-(R)- α -methylbenzylamine (33.4 g, 158 mmol) in THF (450 mL) at 0°C was added n-butyllithium (1.6M in hexanes; 99 mL, 158 mmol). The dark violet solution was stirred at 0°C for 30 minutes, cooled to -78°C, and the ester 3-2 (29.2 g, 150 mmol) in
 25 THF (100 mL) was added over 5 minutes. The resulting solution was stirred at -78°C for 1 hr., then warmed to room temperature. After 2 hrs, the mixture was poured into water and extracted with EtOAc, washed with water, then brine, dried and concentrated *in vacuo* to give

an oil. Column chromatography (silica gel; hexane/EtOAc 1:1 then pure EtOAc) gave the title compound 3-3.

¹H NMR (300 MHz, CDCl₃) δ 1.06 (3H, t), 1.28 (3H, d), 2.52 (1H, dd), 2.62 (1H, dd), 3.66 (1H, d), 3.72 (1H, d), 3.95 (2H, q), 4.44 (1H, dd), 6.95 (1H, m), 7.1-7.5 (13H, m).

3(S)-Fluorophenyl-β-alanine ethyl ester hydrochloride (3-4)

A solution of the N-benzyl-(R)-α-methylbenzylamine 3-3 (28.2 g, 69.6 mmol) in ethanol (300 mL), acetic acid (30 mL) and water (3 mL) was degassed with argon for 30 minutes. Pd(OH)₂ on carbon (20% dry weight; 2.6 g) was added and the mixture then stirred under a hydrogen atmosphere (balloon) for 2 hours. The mixture was filtered through celite and the solvent removed *in vacuo* to give an oil. This oil was dissolved in 200 mL ether, and to this solution was added 60 mL 1N HCl in ether to yield a precipitate. Filtration and washing the solid with ether/hexane gave the title compound 3-4 as a white solid.

¹H NMR (300 MHz, CD₃OD) δ 1.21 (3H, t), 3.0-3.2 (2H, m), 4.16 (2H, q), 4.76 (1H, t), 7.2-7.35 (3H, m), 7.5 (1H, m).


3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (3-5)


Compound 3-5 was prepared from 3-4 using the procedure for the preparation of 2-13.

TLC R_f = 0.36 (15:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

¹H NMR (300 MHz, CD₃OD) δ 7.41 (m, 2H), 7.09 (m, 3H), 6.54 (d, 1H, J=8.2 Hz), 5.48 (m, 1H), 3.51 (m, 2H), 3.46 (m, 3H), 3.23 (m, 2H), 2.94 (m, 2H), 2.81 (m, 4H), 2.63 (m, 2H), 1.93 (m, 2H), 1.19 (m, 2H, J=5.1 Hz).

SCHEME 4





3-Quinolin-3-yl-propionic acid (4-2).

A solution containing quinoline-3-carboxaldehyde 4-1 (5 g, 31.8 mmol), malonic acid (3.6 g, 35.0 mmol), and ammonium acetate (5.0 g, 63.6 mmol) in anhydrous ethanol (125 mL) was heated at reflux for 12 h. After cooling to room temperature, the resulting white solid was collected by filtration and washed with cold ethanol (150 mL) and then dried under vacuum to provide 4-2 as a white solid (3.84 g, 17.8 mmol, 56%).

¹H NMR (300 MHz, D₂O): δ 8.91 (d, J = 2 Hz, 1H), 8.21 (d, J = 2 Hz, 1H), 8.12 (d, J = 8 Hz, 1H), 7.84 (d, J = 7 Hz, 1H), 7.72 (t, J = 7 Hz, 1H), 7.54 (t, J = 7 Hz, 1H), 4.72 (m, 1H), 2.73 (m, 2H).

3-Phenylacetyl-amino-3-quinolin-3-yl-propionic (4-3)

A 0°C solution of 4-2 (3.5 g, 16.2 mmol) and NaHCO₃ (2.7 g, 32.4 mmol) in 50% aqueous dioxane (100 mL) was treated dropwise with a solution of phenylacetyl chloride (3.00 g, 19.4 mmol) in 25 mL of dioxane. The resulting solution was stirred at 0°C for 2.5h., then warmed to room temperature, diluted with H₂O (50 mL) and washed with ether (2 x 100 mL). The aqueous layer was adjusted to pH = 3 with 3N HCl and then extracted with CH₂Cl₂ (3 x 150 mL). The pooled organic extracts were dried, filtered and concentrated to afford 4-3 as an off-white solid.

¹H NMR (300 MHz, CD₃OD): δ 8.85 (d, J = 2 Hz, 1H), 8.20 (d, J = 2 Hz, 1H), 8.00 (d, J = 8 Hz, 1H), 7.86 (d, J = 7 Hz, 1H), 7.76 (t, J = 7 Hz, 1H), 7.52 (t, J

= 7 Hz, 1H), 7.28 (m, 6H), 5.53 (t, J = 6.8 Hz, 1H), 3.57 (s, 2H), 2.96 (m, 2H).

3-(S)-Quinolin-3-yl-propionic acid dihydrochloride (4-6)

5 Acid 4-3 (5.0 g, 15 mmol) was suspended in water (3.5 L), then treated with 1N NaOH (15 mL) to afford a clear solution. Penicillin amidase (Sigma, EC 3.5.1.11, 10,000 U) in 0.1 M phosphate buffer was added. The pH of the mixture was adjusted to 7.8 with 1N NaOH, and the solution was stirred at room temperature for 4 days. The reaction
10 was monitored periodically by HPLC and the reaction stopped once the 50% conversion was reached. Next, the reaction solution was cooled to 0°C and adjusted to pH = 3 with 3N HCl. An oily yellow precipitate formed and was collected by filtration then washed with water to afford crude 4-5 (1.8 g, 5.3 mmol). The filtrate was extracted with CH₂Cl₂ (3 x
15 500 mL) to afford additional 4-5 contaminated by phenylacetic acid. Both batches of crude 4-5 were combined and stirred in 3 N HCl (200 mL) at 50° for 12 h., then cooled, washed with ether (2 x 100 mL) and evaporated to afford 4-6.

20 3-(S)-Quinolin-3-yl-propionic acid ethyl ester dihydrochloride (4-7).


 The resolved acid 4-6 was converted to 4-7 by refluxing in ethanolic HCl.

¹H NMR (300 MHz CD₃OD): δ 9.25 (d, J = 2 Hz 1H), 8.31 (d, J = 2 Hz, 1H), 8.15 (d, J = 8 Hz, 1H), 7.84 (d, J = 7 Hz, 1H), 7.72 (t, J = 7 Hz, 1H), 7.54 (t, J
25 = 7 Hz, 1H), 4.72 (m, 1H), 4.15 (q, J = 6 Hz, 2H), 2.73 (m, 2H) 1.18 (t, J = 6 Hz, 3H).

3(S)-(Quinolin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (4-8).

30 Compound 4-8 was prepared from 4-7 using the procedure for the preparation of 2-13.

¹H NMR (300 MHz, CD₃OD) δ 8.83 (m, 1H), 8.32 (m, 1H), 8.02 (m, 2H), 7.78 (m, 1H), 7.63 (m, 1H), 7.43 (d, 1H, J=7.3 Hz), 6.57 (d, 1H, J=7.3 Hz), 5.76 (m, 1H), 3.73 (q, 1H, J=8.2 Hz), 3.48 (m, 3H), 3.32 (m, 4H), 3.17 (m,
35 2H), 2.95 (m, 1H), 2.84-2.62 (, 6H), 2.10 (m, 1H), 1.88 (m, 3H).

SCHEME 5


5 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid (5-2)

Compound 5-2 was prepared from 5-1 (for preparation, see Zablocki et al., J. Med. Chem. 1995, 38, 2378) using the procedure for the preparation of 2-13.




10 ¹H NMR (300 MHz, CD₃OD) δ 8.54 (m, 1H), 8.47 (m, 1H), 7.85 (d, 1H, J=7.9 Hz), 7.46 (m, 2H), 6.55 (d, 1H, J=7.3 Hz), 5.57 (m, 1H), 3.63 (m, 2H), 3.46 (m, 3H), 3.18 (m, 2H), 3.01 (m, 2H), 2.77 (m, 4H), 2.60 (m, 2H), 2.05 (m, 1H), 1.93 (m, 3H).


TLC (silica): R_f=0.09 (15 EtOAc/10 EtOH/1 NH₄OH/ H₂O)

SCHEME 6



SCHEME 6 (CONTINUED)




SCHEME 6 (CONTINUED)



The imidazolidine-ring methylated compound 6-13 was prepared using the procedure described in Scheme 2 by replacing the glycine ethyl ester with alanine ethyl ester.

In further embodiments, other imidazolidine-ring substituted systems are prepared using the procedure described in Scheme 2 by employing the desired naturally-occurring or non-naturally-occurring amino acid.


10

SCHEME 7


$\text{Pd}(\text{OAc})_2$, Et_3N ,
tri-o-tolylphosphine
ethyl acrylate, CH_3CN











THF
2. $n\text{BuLi}$, THF
3. NH_4Cl , H_2O



$\text{Pd}(\text{OH})_2$, H_2 ,
 EtOH , HOAc , H_2O





5-Bromo-2-methoxypyridine (7-2)

To a solution of KOH (4.2 g, 0.075 mol) in water (750 mL) was added 2-methoxypyridine 7-1 (16.4 g, 0.15 mol) followed by a dropwise addition of bromine (24 g, 0.15 mol) in 1N aqueous KBr (750 mL), and the resulting solution was stirred at room temperature for 5 hr. Solid NaHCO₃ was added until basic, and the solution was extracted with CHCl₃ (3x500 mL). The organic layer was washed with 10% NaHSO₃, then brine, dried over Na₂SO₄, filtered and the solvent removed in vacuo. The resulting dark brown oil was predominantly the desired compound 7-2 and was used as such in the next step.

¹H NMR (300 MHz, CDCl₃) δ 3.91 (3H, s), 6.66 (1H, d), 7.62 (1H, dd), 8.20 (1H, d).

Ethyl 3-(6-methoxypyridin-3-yl)acrylate (7-3)

A solution of the 5-bromo-2-methoxypyridine 7-2 (74.3 g, 0.4 mol), ethyl acrylate (150 mL, 1.4 mol), triethylamine (150 mL, 1.08 mol), palladium acetate (10 g, 0.045 mol) and tri-*o*-tolylphosphine (20 g, 0.066 mol) in 100 mL acetonitrile was degassed with argon for 10 minutes. The mixture is heated at 90°C for 12 hr, then the volatiles were removed in vacuo. Toluene (300 mL) was added and the mixture concentrated again. Diethyl ether (300 mL) was added and the mixture filtered through a pad of silica gel eluting with 800 mL of diethyl ether. After removal of the diethyl ether, the residue was chromatographed on silica gel eluting with EtOAc/hexane 1:19 then 1:14 then 1:9 to give 7-3 as a yellow solid.

¹H NMR (300 MHz, CDCl₃) δ 1.34 (3H, t), 3.97 (3H, s), 4.26 (2H, q), 6.34 (1H, d), 6.76 (1H, d), 7.63 (1H, d), 7.77 (1H, dd), 8.27 (1H, d).

N-Benzyl-(R)-α-methylbenzyl-3(S)-(6-methoxypyridin-3-yl)-β-alanine ethyl ester (7-4)

To a solution of N-benzyl-(R)-α-methylbenzylamine (97.5 g, 462 mmol) in THF (750 mL) at 0°C was added n-butyllithium (2.5M in hexanes; 178.5 mL, 446 mmol). The dark violet solution was stirred at 0°C for 20 minutes, cooled to -78°C and the ester

7-3 (63.7 g, 308 mmol) in THF (250 mL) was added over 60 minutes. The resulting solution was stirred at -78°C for 1 hr., then cannulated into saturated NH₄Cl and extracted with EtOAc, washed with water then brine, dried and concentrated *in vacuo* to give an oil. Column

5 chromatography (silica gel; hexane/EtOAc 9:1 then 4:1) gave 7-4 as an oil contaminated with N-benzyl-(R)- α -methylbenzylamine. This oil was taken up in 5% AcOH in water and extracted with diethyl ether (4x). The organic layers were dried over MgSO₄ and the solvent removed to give the title compound 7-4.

10 ¹H NMR (300 MHz, CDCl₃) δ 1.08 (3H, t), 1.27 (3H, d), 2.52 (1H, dd), 2.62 (1H, dd), 3.66 (1H, d), 3.70 (1H, d), 3.93 (3H, s), 3.95 (2H, m), 4.41 (1H, dd), 6.74 (1H, d), 7.15-7.45 (10H, m), 7.64 (1H, dd), 8.15 (1H, d).

3(S)-(6-methoxypyridin-3-yl)- β -alanine ethyl ester (7-5)

15 To a degassed (argon) solution of the ester 7-4 (70 g) in EtOH (250 mL), HOAc (25 mL) and water (2 mL) was added 20% Pd(OH)₂ on carbon. The mixture was placed under hydrogen using a balloon and the resulting mixture was stirred for 24 hr. After filtration through celite (washing with EtOAc), the solvent was removed in vacuo to afford

20 a waxy solid. This was dissolved in 200 mL water and extracted with diethyl ether (2x200 mL). The aqueous layer was then treated with solid K₂CO₃ until fully saturated and extracted with EtOAc (4x200 mL). After drying over MgSO₄, the solvent was removed in vacuo to give the title compound 7-5 as an oil which solidified in the freezer.

25 ¹H NMR (300 MHz, CDCl₃) δ 1.23 (3H, t), 2.61 (1H, dd), 2.68 (1H, dd), 3.92 (3H, s), 4.15 (2H, q), 4.41 (1H, dd), 6.93 (1H, d), 7.62 (1H, dd), 8.13 (1H, d).

3(S)-(2-(tert-Butoxycarbonyl)-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-amino)-ethylamino)-3-(2-methoxypyridin-5-yl)-propionic acid ethyl ester (7-6)

30 To a solution of the amine 7-5 (6.85 g, 30.5 mmol) in 2-propanol (300 mL) at room temperature was added acetic acid (1.75 mL, 30.5 mmol), NaOAc (24.6 g, 0.3 mol) and 4Å molecular sieves (5 g). The aldehyde 2-9A (8.1 g, 24.3 mmol) in 2-propanol (150 mL) was added and

the mixture stirred for 15 minutes, then cooled to 0°C and NaCNBH₃ (5.66 g, 90 mmol) added in one lot. The resulting mixture was allowed to warm to room temperature and stirred for 16 hr before being filtered through celite. After removal of the solvent in vacuo, the residue was treated with 10% aqueous KHSO₄ for 30 minutes, basified with solid K₂CO₃ (to pH ~10) and extracted with 3x 200 mL EtOAc. The EtOAc layers were washed with brine, dried (Na₂SO₄) and concentrated *in vacuo* to give an oil. Column chromatography (silica gel; 5% MeOH in CHCl₃) gave 7-6 as an oil contaminated with ~7% of the β-alanine.

¹H NMR (300 MHz, CDCl₃) δ 1.20 (3H, t), 1.42 (9H, s), 1.7-2 (4H, br m), 2.5-2.8 (8H, m), 3.2 (4H, m), 3.42 (2H, m), 3.92 (3H, s), 4.06 (2H, q), 5.0-5.4 (1H, bs), 6.36 (1H, br s), 6.72 (1H, d), 7.12 (1H, br s), 7.58 (1H, dd), 8.07 (1H, d).

15 3(S)-(6-Methoxypyridin-3-yl)-3-(2-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propylaminol]-ethylaminol)-propionic acid ethyl ester (7-7)

A solution of the ester 7-6 (14.1 g, 26 mmol) in EtOAc (350 mL) cooled to -20°C was treated with HCl (gas) for 10 minutes then stoppered and stirred at 0°C for 1.5 hr. The volatiles were removed in vacuo, the residue taken up in 150 mL of water and treated with solid K₂CO₃ to pH~10. This solution was extracted with EtOAc (3x 150 mL), washed with brine, dried (Na₂SO₄) and concentrated to give an oil. Column chromatography (silica gel; 5% MeOH in CHCl₃) gave the β-alanine 7-5; further elution with 5% MeOH in CHCl₃ saturated with NH₃ gave 7-7 as a viscous oil.

¹H NMR (300 MHz, CDCl₃) δ 1.20 (3H, t), 1.8-1.95 (4H, m), 2.5-2.8 (12H, m), 3.39 (2H, m), 3.92 (3H, s), 4.09 (2H, q), 5.01 (1H, bs), 6.34 (1H, d), 6.72 (1H, d), 7.06 (1H, d), 7.59 (1H, dd), 8.07 (1H, d).

30 3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid ethyl ester (7-8)

To a solution of the diamine 7-7 (8.03 g, 18.2 mmol), DIPEA (9.5 mL, 54.6 mmol) and DMAP (250 mg) in 1,2-dichloroethane (150 mL)

cooled to -20°C was added p-nitrophenyl chloroformate (3.85 g, 19.1 mmol) in 1,2-dichloroethane (25 mL) dropwise such that the internal temperature remains below -15°C. The resulting mixture was allowed to warm to 0°C and stirred for 45 minutes, then heated to reflux for 4 hr.

5 After cooling, the solvent was evaporated in vacuo, the residue taken up in EtOAc and washed successively with 10% K₂CO₃ (6x 150 mL) and brine. The EtOAc layer was dried (Na₂SO₄) and concentrated *in vacuo* to give an oil (6.27 g). Column chromatography (silica gel; 5% EtOH in CH₂Cl₂) gave 7-8 as an oil.

10 ¹H NMR (300 MHz, CDCl₃) δ 1.20 (3H, t), 1.8-1.95 (4H, m), 2.52 (2H, dd), 2.68 (1H, dd), 2.9-3.1 (3H, m), 3.15-3.3 (5H, m), 3.39 (2H, m), 3.92 (3H, s), 4.11 (2H, q), 4.8 (1H, bs), 5.42 (1H, t), 6.34 (1H, d), 6.72 (1H, d), 7.03 (1H, d), 7.60 (1H, dd), 8.08 (1H, d).

15 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (7-9)


To a solution of the ester 7-8 (3.48 g, 7.4 mmol) in MeOH (50 mL) and water (30 mL) at room temperature was added 1N NaOH solution (22.3 mL, 22.3 mmol) and the mixture stirred for 16 hr. After
20 removal of the solvent in vacuo, the residue was treated with 25 mL 1N HCl and the solvent removed again. Column chromatography of the residue (silica gel; EtOAc/EtOH/aq. NH₄OH/H₂O 20:10:1:1 then 15:10:1:1) gave a gum which was crystallized from a minimum amount of water and filtered to give 7-9 as a white solid.

25 ¹H NMR (300 MHz, CD₃OD) δ 1.75-2.1 (4H, m), 2.55-3.1 (8H, m), 3.28 (1H, q), 3.3 (1H, m), 3.4-3.55 (3H, m), 3.63 (1H, q), 3.85 (3H, s), 5.47 (1H, dd), 6.55 (1H, d), 6.80 (1H, d), 7.48 (1H, d), 7.68 (1H, d), 8.09 (1H, d).

In a further embodiment, the R-enantiomer, i.e. 3(R)-(6-methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid was prepared by substituting N-benzyl-(S)-α-methylbenzylamine for the N-benzyl-(R)-α-methylbenzylamine in preparing intermediate 7-4.
30

In yet a further embodiment, the racemate, i.e. 3-(6-methoxypyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid was prepared by substituting racemic N-benzyl- α -methylbenzylamine for the N-benzyl-
 5 (R)- α -methylbenzylamine in preparing intermediate 7-4.

SCHEME 8



5-Bromo-2-ethoxypyridine (8-2)

10 Sodium metal (4.87 g, 0.212 mol) was added to ethanol (200 mL) and stirred until completely dissolved. To this solution was added 2,5-dibromopyridine 8-1 (Aldrich; 10 g, 0.0424 mol) and the resulting mixture was stirred at reflux for 16 hr. The solvent was removed in vacuo and the residue partitioned between water and EtOAc. After
 15 extraction with EtOAc (2x), the organic layer was washed with brine, dried (MgSO₄) and concentrated to give 8-2 as a red-brown solid which was used as such in the next step.

¹H NMR (300 MHz, CDCl₃): δ 1.4 (3H, t), 4.33 (2H, q), 6.63 (1H, d), 7.62 (1H, dd), 8.19 (1H, d).

20

3(S)-(6-Ethoxypyridin-3-yl)- β -alanine ethyl ester (8-3)

The title compound 8-3 was prepared from 8-2 using the procedure described for the synthesis of 7-5.


¹H NMR (300 MHz, CDCl₃): δ 1.25 (3H, t), 1.39 (3H, t), 2.61 (1H, dd), 2.67
5 (1H, dd), 4.15 (2H, q), 4.34 (2H, q), 4.40 (1H, dd), 6.71 (1H, d), 7.62 (1H, dd),
8.11 (1H, d).

3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid (8-4)

10 The title compound 8-4 was prepared from 2-9A and 8-3 using the procedure described in Scheme 2.

¹H NMR (300 MHz, CD₃OD) δ 1.37 (3H, t), 1.8-2.0 (4H, m), 2.65 (2H, t),
2.82 (2H, t), 2.95-3.10 (2H, m), 3.15 (2H, m), 3.23 (2H, dt), 3.46 (2H, m), 3.51
(2H, t), 4.32 (2H, q), 5.41 (1H, t), 6.62 (1H, d), 6.84 (1H, d), 7.57 (1H, d), 7.76
15 (1H, dd), 8.13 (1H, d).

SCHEME 9

3-(6-Amino-pyridin-3-yl)-acrylic acid *tert*-butyl ester (9-2)

A mixture of 2-amino-5-bromo-pyridine (**9-1**) (10 g, 58 mmol), *tert*-butyl acrylate (50 mL, 344 mmol), triethylamine (50 mL, 359 mmol), tri-*o*-tolylphosphine (3.0 g, 9.8 mmol) and $\text{Pd}(\text{OAc})_2$ (1.0 g, 4.5 mmol) in 150 mL CH_3CN was purged with argon for 5 min and subsequently refluxed at 110°C for 20 hr. The mixture was then cooled and concentrated. The residue was purified using silica gel flash

chromatography (EtOAc/hexanes 1:1) to afford the desired product 9-2 as a solid.

R_f (silica, EtOAc/hexanes 1:1) = 0.26

5 3(S)-(6-Amino-pyridin-3-yl)-3-[benzyl-(1(R)-phenylethyl)-amino]-propionic acid *tert*-butyl ester (9-3)

To a cooled (0°C) solution of (R)-(+)-N-benzyl- α -methylbenzylamine (4.0 g, 19 mmol) in 50 mL THF was gradually added n-butyllithium (11.3 mL, 2.5 M, 28.2 mmol) over 5 min. The mixture was stirred for 30 min at 0°C and cooled to -78°C. A solution of 9-2 (2.0 g, 9.4 mmol) in 20 mL THF was gradually added. After stirring for 40 min at -78°C, it was treated with NH₄Cl (sat.) at -78°C, warmed to room temperature and extracted three times with EtOAc. The combined organic layers were washed with brine and dried over Na₂SO₄. After solvent evaporation, the residue was purified using silica gel flash chromatography (EtOAc/hexanes, 1:2) to afford the desired product 9-3 as an oil.

R_f (silica, EtOAc/hexanes 1:1) = 0.28

20 3(S)-Amino-3-(6-amino-pyridin-3-yl)-propionic acid ethyl ester•2 HCl (9-4)

A mixture of 9-3 (0.5 g, 1.2 mmol) and 10% Pd/C (0.4 g) in 10 mL AcOH was purged with argon for 5 min and then heated at 78°C. 1,4-Cyclohexadiene (2 mL, 21.1 mmol) was then gradually added. The reaction mixture was stirred for 3 hr and filtered through a celite pad. The solution was concentrated and the residue was purified using silica gel flash chromatography (EtOAc/MeOH/NH₄OH 1 :1:0.04) to afford an oil. To the oil (1.2 g) in 20 mL EtOH was introduced HCl gas for 10 min. The mixture was stirred 24 hr and then concentrated to afford the desired product 9-4 as the HCl salt.

30 ¹H NMR (400 MHz, CD₃OD) δ 8.11 (d, J=9.6 Hz, 1H), 8.08 (s, 1H), 7.13 (d, J=9.6 Hz, 1H), 4.77 (m, 1H), 4.18 (q, J=6.8 Hz, 2H), 3.22-3.02 (m, 2H), 1.24 (t, J=6.8 Hz, 3H).


3(S)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid (9-5)


The title compound 9-5 was prepared as the TFA salt from 2-9A and 9-4 using the procedure described in Scheme 2.

- 5 ¹H NMR (300 MHz, CD₃OD) δ 7.90 (d, J=2.1 Hz, 1H), 7.46 (dd, J=8.7, 2.1 Hz, 1H), 7.03 (d, J=8.7 Hz, 1H), 6.47 (d, J=8.7 Hz, 1H), 6.34 (d, J=8.7 Hz, 1H), 5.38-5.30 (m, 1H), 3.40-3.37 (m, 2H), 3.26-3.16 (m, 4H), 3.03-2.86 (m, 2H), 2.70-2.66 (m, 2H), 2.55-2.50 (m, 2H), 2.14-2.02 (m, 2H), 1.93-1.79 (m, 4H).

10

SCHEME 10





3-(3-Hydroxy-4-nitrophenyl)-acrylic acid ethyl ester (10-2)

To a stirred solution of aldehyde 10-1 (20.28 g, 132.5 mmol) in
 5 CH_2Cl_2 (400 mL) at room temperature was added
 (carbethoxymethylene)triphenylphosphorane (46.12 g, 132.5 mmol) over
 a 10 min period. The resulting orange solution was stirred at room
 temperature for 2 h. The solution was concentrated to one-fourth its
 volume. Flash chromatography (silica gel; 30:70 EtOAc/hexanes) gave
 10 the title compound 10-2 as a bright yellow solid.
 TLC $R_f = 0.75$ (25:75 EtOAc/hexanes)
 ^1H NMR (300 MHz, CDCl_3) δ 8.14 (d, 1H), 7.60 (d, 1H), 7.15 (dd, 1H), 6.54
 (d, 1H), 4.30 (q, 2H), 1.36 (t, 3H).

3-(4-Amino-3-hydroxyphenyl)-acrylic acid ethyl ester (10-3)

To a stirred suspension of 10-2 (4.64 g, 19.6 mmol), NH₄Cl (524 mg, 9.8 mmol), EtOH (140 mL) and H₂O (70 mL) was added iron dust (2.72 g, 48.9 mmol). The resulting yellow suspension was refluxed for 1.5 h. and then the solution was filtered while hot through celite. The filtrate was concentrated and the residue was partitioned between EtOAc and brine. The layers were separated and the EtOAc layer dried (Na₂SO₄) and concentrated to give 10-3 which was used without further purification in the next step.

TLC R_f = 0.2 (25:75 EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 7.57 (d, 1H), 7.00 (m, 2H), 6.68 (d, 1H), 6.20 (d, 1H), 4.26 (q, 2H), 4.10 (bs, 2H), 1.33 (t, 3H).

3-[4-(2-Chloroacetyl-amino)-3-hydroxyphenyl]acrylic acid ethyl ester (10-4)

To a stirred solution of 10-3 (3.38 g, 16.3 mmol) in CHCl₃ (80 mL) was added saturated NaHCO₃ (50 mL) and it was then chilled to 0°C. A solution of chloroacetyl chloride (1.94 mL, 24.4 mmol) in CHCl₃ (30 mL) was added dropwise to the chilled biphasic. Upon addition completion, the reaction was stirred at 0°C for 1 h. The layers were separated and the aqueous layer was extracted twice with EtOAc. The combined organic layers were washed with brine, dried (Na₂SO₄) and concentrated to give 10-4 which was used without further purification in the next step.

TLC R_f = 0.4 (25:75 EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃): δ 10.33 (s, 1H), 9.58 (s, 1H), 8.02 (d, 1H), 7.51 (d, 1H), 7.19 (d, 1H), 7.12 (s, 1H), 6.39 (d, 1H), 4.42 (s, 2H), 4.17 (q, 2H), 1.25 (t, 3H).

3-(3-Oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl) acrylic acid ethyl ester (10-5)

To a stirred solution of 10-4 (4.28 g, 15.0 mmol) in DMF (50 mL) was added K₂CO₃ (4.50 g, 32.6 mmol). The resulting suspension was heated to 50°C for 12 h., after which time the reaction was concentrated. The residue was partitioned between saturated NaHCO₃

and EtOAc and extracted twice with EtOAc. The combined organic layers were washed with brine, dried (Na₂SO₄), and concentrated.

Flash chromatography (silica gel; 25:75 EtOAc/hexanes) yielded 10-5 as a beige solid.

5 TLC Rf = 0.5 (25:75 EtOAc/hexanes).

¹H NMR (300 MHz, CDCl₃) δ 10.91 (s, 1H), 7.54 (d, 1H), 7.37 (s, 1H), 7.31 (d, 1H), 6.90 (d, 1H), 6.51 (d, 1H), 4.60 (s, 2H), 4.16 (q, 2H), 1.24 (t, 3H).

10 3(R)-[Benzyl-(1-phenylethyl)-aminol-3-(S)-(3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl) propionic acid ethyl ester (10-6)

To a stirred solution of (R)-(+)-N-benzyl-α-methylbenzylamine (5.43 g, 25.7 mmol) and anhydrous THF (75 mL) at 0°C was added butyllithium (10.3 mL, 2.5 M/hexanes, 25.7 mmol) via syringe. The violet-red solution was stirred at 0°C for 15 minutes and
15 then cooled to -78°C. A solution of 10-5 (2.12 g, 8.6 mmol) in anhydrous THF (50 mL) was added via syringe, and the resulting brown solution was stirred at -78°C for 30 minutes. The brown solution was quenched with saturated NH₄Cl, the mixture then warmed to room temperature and extracted twice with Et₂O. The combined organic layers were
20 washed with brine, dried (Na₂SO₄), and concentrated. Flash chromatography (silica gel; 15:85 to 25:75 EtOAc/hexanes) yielded 10-6 as a white foam.

TLC Rf = 0.25 (25:75 EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 10.89 (s, 1H), 7.32 (m, 10H), 7.10 (m, 2H),
25 6.91 (d, 1H), 4.62 (s, 2H), 4.39 (m, 1H), 4.13 (q, 2H) 3.96 (m, 1H), 3.68 (s, 2H), 2.56 (m, 2H), 1.28 (m, 6H).

3(R)-[Benzyl-(1-phenylethyl)-aminol-3-(S)-(4-methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl) propionic acid ethyl ester (10-7)

30 To a stirred suspension of NaH (65 mg, 60%, 1.6 mmol) in DMF (5 mL) under argon was added a solution of 10-6 (650 mg, 1.4 mmol) in DMF (10 mL) via syringe. This yellow solution was stirred at room temperature for 30 minutes. Iodomethane (0.5 mL, 8.0 mmol) was added and the solution then stirred at room temperature for an
35 additional 30 minutes. The reaction was quenched with saturated

NaHCO₃. The aqueous layer was extracted three times with CH₂Cl₂. The combined organic layers were washed with brine, dried (Na₂SO₄) and concentrated. Flash chromatography (silica gel; 25:75 EtOAc/hexanes) afforded 10-7 as a clear oil.

- 5 TLC Rf = 0.6 (25:75 EtOAc/hexanes)
1H NMR (300 MHz, CDCl₃) δ 7.30 (m, 10H), 7.06 (m, 2H), 6.91 (d, 1H), 4.62 (s, 2H), 4.39 (m, 1H), 4.13 (q, 2H) 3.96 (m, 1H), 3.68 (s, 2H), 3.35 (s, 3H), 2.56 (m, 2H), 1.26 (m, 6H).

- 10 3(S)-Amino-3-(4-methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)propionic acid ethyl ester (10-8)

A stirred solution of 10-7 (581 mg, 1.2 mmol), MeOH (10 mL), AcOH (1.0 mL), and H₂O (0.3 mL) was degassed with argon for 5 minutes. Pd(OH)₂ (581 mg) was added and the reaction was placed
15 under 1 atm of H₂ for 2.5 h. The reaction was diluted with EtOAc and filtered through celite. The filtrate was concentrated to yield 10-8 as a clear oil.

- TLC Rf = 0.3 (5:95 MeOH/CH₂Cl₂)
1H NMR (300 MHz, CDCl₃) δ 7.04 (m, 2H), 6.93 (dd, 1H), 4.61 (s, 2H), 4.39
20 (m, 1H), 4.13 (q, 2H), 3.37 (b, 2H), 3.35 (s, 3H), 2.69 (m, 2H), 1.24 (t, 3H).


3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid (10-9)

- 25 The title compound 10-9 was prepared from 2-9A and 10-8 using the procedure described in Scheme 2.

1H NMR (400 MHz, d₆-DMSO) δ 7.83 (bs, 1H), 7.60 (d, 1H), 7.11 (d, 1H), 6.99 (d, 1H), 6.93 (s, 1H), 6.63 (d, 1H), 5.20 (t, 1H), 4.64 (s, 2H), 3.3-2.8 (m, 10H), 3.25 (s, 3H), 2.72 (m, 2H), 2.59 (m, 2H), 1.81 (m, 2H), 1.74 (m, 2H).

30

SCHEME 11



2-tert-Butoxycarbonylamino-5-aminopyridine (11-2)

A solution of 2-amino-4-bromopyridine 11-1 (10.1 g, 58.4 mmol) in 150 mL of melted *t*-BuOH was treated with di-*tert*-butyl dicarbonate (14.0 g, 64.2 mmol). After the solution was stirred for 12 hr, the solvent was evaporated. The residue was purified using silica gel flash chromatography (CHCl₃/hexanes, 5:1) to afford the desired product 11-2 as a solid.

R_f(silica, 100% CHCl₃) = 0.56

¹H NMR (300 MHz, CDCl₃) δ 8.82 (bs, 1H), 8.38 (d, 1H), 8.78 (d, 1H), 7.78 (dd, 1H), 1.55 (s, 9H).

2-(tert-Butoxycarbonyl-methyl-amino)-5-aminopyridine (11-3)

To a solution of 11-2 (6.0 g, 22.0 mmol) in 50 mL DMF at 0°C was added NaH gradually. After the mixture was stirred for 40 min, CH₃I (3.4 g, 24.0 mmol) was added in one portion. The reaction mixture was stirred for 5 hr, treated with 300 mL water and extracted three times with ethyl ether. The combined organic layers were washed with brine and dried over Na₂SO₄. After solvent removal, the residue was purified by silica gel flash chromatography (CHCl₃/hexanes 6:1) to afford the desired product 11-3 as a solid.

R_f(silica, 100% CHCl₃) = 0.40

¹H NMR (300 MHz, CDCl₃) δ 8.40 (dd, 1H), 7.68 (m, 2H), 3.36 (s, 3H), 1.55 (s, 9H).

3-[6-(tert-Butoxycarbonyl-methyl-amino)-pyridin-3-yl]-acrylic acid ethyl ester (11-4)

A mixture of 11-3 (6.0 g, 20.9 mmol), ethyl acrylate (6.3 mL, 62.7 mmol), triethylamine (17 mL, 125.5 mmol), tri-*o*-tolylphosphine (1.3 g, 6.2 mmol) and Pd(OAc)₂ (0.5 g, 2.1 mmol) in 50 mL CH₃CN was purged with argon for 5 min and subsequently refluxed at 110°C for 20 hr. The mixture was cooled and concentrated. The residue was purified using silica gel flash chromatography (EtOAc/hexanes 1:3) to afford the desired product 11-4 as an oil.

¹H NMR (300 MHz, CDCl₃) δ 8.47 (bs, 1H), 7.82 (m, 2H), 7.64 (d, 1H), 6.42 (d, 1H), 4.27 (q, 2H), 3.43 (s, 3H), 1.54 (s, 9H), 1.34 (t, 3H).

3-Benzylamino-3-[6-(tert-butoxycarbonyl-methyl-amino)-pyridin-3-yl]-propionic acid ethyl ester (11-5)

A mixture of 11-4 (1.7 g, 5.6 mmol) and benzylamine (8 mL, 73.2 mmol) was heated in a sealed-tube at 95°C for 24 hr. The crude reaction mixture was purified using silica gel flash chromatography (EtOAc/hexanes 1:3 to 1:1) to afford the desired product 11-5 as an oil. R_f (silica, EtOAc/hexanes 1:1) = 0.63.


10 3-Amino-3-[6-(tert-butoxycarbonyl-methyl-amino)-pyridin-3-yl]-propionic acid ethyl ester (11-6)

A mixture of 11-5 (1.5 g 3.6 mmol), 20% Pd(OH)₂/C (0.3 g), AcOH (5.5 mL) and EtOH (50 mL) was purged with argon 3 times under vacuum. The reaction mixture was stirred under balloon hydrogenation condition for 16 hr and filtered through a celite pad. After solvent removal, the desired product 11-6 was obtained as the acetate salt. ¹H NMR (300 MHz, CDCl₃) δ 8.38 (d, 1H), 7.70 (m, 2H), 4.50 (dd, 1H), 4.15 (q, 2H), 3.40 (s, 3H), 2.80 (m, 2H), 1.25 (t, 3H).

20 3-(6-Methylamino-pyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (11-7)

The title compound 11-7 was prepared as the TFA salt from 2-9A and 11-6 using the procedure described in Scheme 2. ¹H NMR (300 MHz, CD₃OD) δ 7.92 (dd, J=9.6, 1.6 Hz, 1H), 7.79 (d, J=1.6 Hz, 1H), 7.57 (d, J=7.6 Hz, 1H), 7.03 (d, J=9.6 Hz, 1H), 6.63 (d, J=7.6 Hz, 1H), 5.28 (m, 1H), 3.51-3.36 (m, 5H), 3.28-3.17 (m, 3H), 3.05 (m, 2H), 3.02 (s, 3H), 2.82 (m, 2H), 2.67 (m, 2H), 1.98-1.84 (m, 4H).

SCHEME 12

3-(2-Fluoro-biphenyl-4-yl)-acrylic acid ethyl ester (12-2)

A solution of 2-fluoro-4-bromobiphenyl **12-1** (7.5 gm, 31.8 mmol), ethyl acrylate (4.3 mL), $\text{Pd}(\text{OAc})_2$ (0.714 gm, 3.2 mmol), tri-*o*-tolylphosphine (1.94 gm, 1.5 mmol), and triethylamine (12 mL) was

heated to 100°C in a sealed tube for 12 h. The reaction was cooled to room temperature and diluted with dichloromethane (40 mL). The organic solution was washed with 10% aq. citric acid (20 mL), satd. aq. NaHCO₃, and brine (20 mL). The organic solution was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (95:5 to 90:10 hexanes/EtOAc) to give the acrylate ester 12-2 as a white solid.

TLC Rf = 0.44 (10% ethyl acetate/hexanes).

10 3-[Benzyl-(1(R)-phenylethyl)-amino]-3-(2-fluoro-biphenyl-4-yl)-propionic acid ethyl ester (12-3)

A cooled (0°C) solution of (R)-(+)-N-benzyl- α -methylbenzylamine (8.9 mL, 42.6 mmol) in THF (100 mL) was treated with n-butyllithium (26.6 mL of a 1.6 M soln in hexanes; 42.6 mmol). After stirring for 10 min, the purple solution was cooled to -78°C and treated with a solution of ester 12-2 (5.76 g, 21.3 mmol) in THF (10 mL). After stirring for 20 min, the solution was quenched with satd aq NH₄Cl soln (5 mL), and the cold bath removed. The reaction mixture was diluted with Et₂O (100 mL), and washed with 10% aq citric acid (50 mL), satd aq NaHCO₃ (50 mL), 5% aq acetic acid (30 mL), 10% aq K₂CO₃ (50 mL), and brine (50 mL). The solution was dried over MgSO₄, filtered and concentrated. The residue was purified by flash chromatography (90:10 hexanes/EtOAc) to give adduct 12-3.

TLC Rf = 0.48 (10% ethyl acetate/hexanes).

25 3-Amino-3-(2-fluoro-biphenyl-4-yl)-propionic acid ethyl ester (12-4)

A solution of the dibenzylamine 12-3 (5.65 gm, 11.75 mmol) in EtOH/HOAc (90/10 mL) was purged with argon and treated with Pd(OH)₂ (3 g) and placed under 1 atm of H₂ gas for 12 h. Additional portions (2.5 g) of Pd(OH)₂ were added after 24 h, 48 h and 144 h. The reaction mixture was purged with argon, filtered through Celite, and the filtrate dissolved in aq HCl (pH=1). The aqueous solution was washed with EtOAc, neutralized with satd aq NaHCO₃, and extracted with EtOAc (3 x 30 mL). The combined organic solutions were washed

with brine, dried over MgSO_4 , filtered and concentrated to give the desired product 12-4.

^1H NMR (300 MHz, CD_3OD) δ 7.41 (m, 8H), 4.10 (m, 1H), 4.06 (m, 2H), 2.73 (m, 2H), 1.18 (m, 3H) ppm.


5

3(S)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl) propionic acid (12-5)

The title compound 12-5 was prepared from 2-9A and 12-4 using the procedure described in Scheme 2.

10 ^1H NMR (300 MHz, CD_3OD) δ 7.49 (m, 9H), 6.64 (d, $J = 7.3$ Hz, 1H), 5.49 (m, 1H), 3.31 (m, 9H), 2.83 (m, 2H), 2.74 (m, 2H), 1.97 (m, 4H) ppm.

SCHEME 13

3-(3-Hydroxy-4-nitro-phenyl)-acrylic acid ethyl ester (13-2)

To a solution of aldehyde 13-1 (15.0 g, 98.0 mmol) in CH_2Cl_2
 5 (300 mL) was slowly added carboethoxymethylenetriphenylphosphorane

(34.1 g, 98.0 mmol). The orange solution was stirred for 12 h at ambient temperature. The solution was concentrated to a paste and purified by flash chromatography (10% EtOAc/CH₂Cl₂) to give 13-2 as a yellow solid. TLC R_f = 0.51 (30% ethyl acetate/hexanes).

- 5 ¹H NMR (300 MHz, CD₃OD) δ 8.08 (d, J=8.4 Hz, 1H), 7.63 (d, J=16.2 Hz, 1H), 7.35 (d, J=1.5 Hz, 1H), 7.27 (dd, J=8.4, 1.5 Hz, 1H), 6.65 (d, J=15.9 Hz, 1H), 4.25 (q, J=7.2 Hz, 2H), 1.32 (t, J=6.9 Hz, 3H) ppm.

3-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-acrylic acid ethyl ester (13-3)

- 10 To a solution of the nitrophenol 13-2 (12.0 g, 57.4 mmol) in warm (70°C) AcOH/H₂O (200 mL) was added iron dust (9.61 g, 172.2 mmol). The brown heterogeneous mixture was stirred for 30 min at 70-80°C. The mixture was filtered hot through Celite, and the Celite bed washed with EtOAc (2 x 200 mL). The filtrate was cautiously neutralized
15 with satd aq NaHCO₃ (3 x 100 mL). The solution was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (5% MeOH in CH₂Cl₂) to give an orange solid (9.6 g, 81%). A portion of this solid (4.5 g, 21.7 mmol) was dissolved in THF (150 mL) and treated with 1,1-carbonyldiimidazole (3.87 g, 23.8 mmol), and
20 the solution was stirred at ambient temperature for 24 h. The solution was diluted with EtOAc (100 mL) and washed with 10% HCl (50 mL) and brine (50 mL). The solution was dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (5% MeOH in CH₂Cl₂) to give 13-3 as a yellow solid.
25 TLC R_f = 0.49 (5% MeOH/CH₂Cl₂).
¹H NMR (300 MHz, CD₃OD) δ 7.77 (d, J=15.9 Hz, 1H), 7.55 (s, 1H), 7.41 (d, J=8.4 Hz, 1H), 7.09 (d, J=8.1 Hz, 1H), 6.47 (d, J=15.9 Hz, 1H), 4.22 (q, J=7.2 Hz, 2H), 1.31 (t, J=7.2 Hz, 3H) ppm.

- 30 3(S)-Amino-3-(2-oxo-2,3-dihydro-benzoxazol-6-yl)-propionic acid ethyl ester (13-4)

- A solution of (R)-(+)-N-benzyl-α-methylbenzylamine (4.08 g, 19.3 mmol) in THF (120 mL) at 0°C was treated with n-BuLi (7.72 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0°C for 30
35 min and then cooled to -78°C. A solution of acrylate 13-3 (1.5 g, 6.43


mmol) in THF (20 mL) was added. After stirring for 15 min at -78°C, satd aq NH₄Cl soln (25 mL) was added and the cold bath removed. The mixture was warmed to room temperature and extracted with Et₂O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL),
5 dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (30% ethyl acetate/hexanes) to give 2.74 g of the β-aminoester as a yellow oil. The aminoester was dissolved in EtOH/H₂O/AcOH (54 mL/4.8 mL/1.2 mL), degassed with argon, and treated with Pd(OH)₂ (2.74 g). The mixture was placed under 1 atm of
10 H₂. After stirring for 18 h, the mixture was diluted with EtOAc and filtered through Celite. The filtrate was concentrated to give ester 13-4 as an off-white solid.
TLC R_f = 0.10 (5% MeOH/CH₂Cl₂).
¹H NMR (300 MHz, CD₃OD) δ 7.34 (s, 1H), 7.26 (dd, J=1.2, 8.1 Hz, 1H),
15 7.12 (d, J=8.1 Hz, 1H), 4.65 (t, J=7.2 Hz, 1H), 4.13 (q, J=6.9 Hz, 2H), 2.98 (m, 2H), 1.20 (t, J=7.2 Hz, 3H) ppm.

3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl} propionic acid (13-5)

20 The title compound 13-5 was prepared from 2-9A and 13-4 using the procedure described in Scheme 2.
¹H NMR (300 MHz, CD₃OD) δ 7.57 (d, J=7.3 Hz, 1H), 7.28 (s, 1H), 7.19 (d, J=8.2 Hz, 1H), 6.63 (d, J= 7.3 Hz, 1H), 5.47 (m, 1H), 3.30 (m, 9H), 2.82 (m, 2H), 2.66 (m, 2H), 1.96 (m, 6H) ppm.

25

SCHEME 14

3-(4-Hydroxy-3-fluorophenyl)-acrylic acid ethyl ester (14-2)

A solution of 2-fluoro-4-bromophenol **14-1** (50 g, 261.8 mmol),
 5 ethyl acrylate (34 mL), $\text{Pd}(\text{OAc})_2$ (2.5 g), tri-*o*-tolylphosphine (5 g) and

triethylamine (83 mL) was heated to 100°C in a sealed tube for 12 h. The reaction was cooled to room temperature and diluted with dichloromethane (100 mL). The organic solution was washed with 10% aq. citric acid (40 mL), satd aq NaHCO₃, and brine (40 mL). The organic
5 solution was dried over MgSO₄, filtered and concentrated. The residue was purified by flash chromatography (50:50 hexanes/EtOAc to 100% EtOAc) to give acrylic acid 14-2 as a white solid.
TLC Rf = 0.45 (50% ethyl acetate/hexanes).

10 3-[Benzyl-(1(R)-phenylethyl)-aminol-3-(4-ethoxy-3-fluorophenyl)-propionic acid ethyl ester (14-4)

To a stirred solution of 14-2 (49.25 gm, 234.5 mmol) in DMF (600 mL) was added Cs₂CO₃ (84.1 gm, 257.9 mmol) and ethyl iodide (18.8 mL, 234.5 mmol). After stirring for 12 h at room temperature, the
15 reaction mixture was diluted with EtOAc (1L) and washed with water (6 x 300 mL), 10% aq. citric acid (200 mL), satd. aq. NaHCO₃ (200 mL), and brine (300 mL). The organic solution was dried over MgSO₄, filtered, and concentrated to give 52.9 g (95%) of the product 14-3 as an orange oil which crystallized upon standing. A cooled (0°C) solution of (R)-(+)-N-benzyl- α -methylbenzylamine (71 mL, 339.4 mmol) in THF (650 mL) was
20 treated with n-butyllithium (212 mL of a 1.6 M soln in hexanes; 339.4 mmol). After stirring for 10 min, the purple solution was cooled to -78°C and treated with a solution of ester 14-3 (53.8 g, 226.3 mmol) in THF (100 mL). After stirring for 20 min, the solution was quenched with satd aq
25 NH₄Cl soln (50 mL), and the cold bath removed. The reaction mixture was diluted with Et₂O (1000 mL), and washed with 10% aq citric acid (300 mL), satd aq NaHCO₃ (300 mL), 5% aq acetic acid (300 mL), 10% aq K₂CO₃ (300 mL), and brine (200 mL). The solution was dried over MgSO₄, filtered and concentrated. The residue was purified by flash
30 chromatography (85:15 hexanes/EtOAc) to give the adduct 14-4.
TLC Rf = 0.39 (25% ethyl acetate/hexanes).

3-Amino-3-(4-Ethoxy-3-fluorophenyl)-propionic acid ethyl ester (14-5)

A solution of the dibenzylamine 14-4 (30.0 gm, 66.8 mmol) in
35 EtOH/HOAc (340/30 mL) was purged with argon and treated with

Pd(OH)₂ (6 g) and placed under 1 atm of H₂ for 12 h. Additional portions (2.5 g) of Pd(OH)₂ were added after 24 h and 48 h. The reaction mixture was purged with argon, filtered through Celite, and the filtrate collected. The filtrate was concentrated to yield the desired amine 14-5.

5 ¹H NMR (300 MHz, CD₃OD) δ 7.19 (m, 3H), 4.62 (m, 1H), 4.07 (m, 4H), 2.99 (m, 2H), 1.39 (m, 3H) 1.18 (m, 3H) ppm.


3-(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl) propionic acid (14-6)

10 The title compound 14-6 was prepared from 2-9A and 14-5 using the procedure described in Scheme 2.

¹H NMR (300 MHz, CD₃OD) δ 7.45 (d, J = 7.3 Hz, 1H), 7.04 (m, 3H), 6.53 (d, J=7.3 Hz, 1H), 5.43 (m, 1H), 4.06 (q, J= 7.0 Hz, 2H), 3.48 (m, 6H), 3.15 (m, 1H), 2.78 (m, 6H), 2.55 (m, 2H), 1.96 (m, 3H), 1.38 (t, J= 7.0 Hz, 3H)

15 ppm.

SCHEME 15



5-Ethoxy-nicotinic acid ethyl ester (15-2)

A mixture of 3-hydroxy-nicotinic acid methyl ester 15-1 (15 g, 90.8 mmol), ethyl iodide (14.5 ml, 181.6 mmol), cesium carbonate (29.5 g, 90.8 mmol) and DMF (150 mL) was stirred at ambient temperature for 3 hours. The reaction mixture was diluted with Et₂O and then washed with 10% K₂CO₃, brine, dried (MgSO₄), and concentrated to give the ester 15-2 as a red oil.

TLC R_f = 0.52 (silica, 75% EtOAc/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 8.82 (s, 1H), 8.46 (s, 1H), 7.75 (s, 1H), 4.40 (q, 2H, J=7Hz), 4.12 (q, 2H, J=7Hz), 1.43 (m, 6H).

5-Ethoxy-N-methoxy-N-methyl-nicotinamide (15-3)

To a solution of 15-2 (15 g, 72 mmol) in EtOH (100 mL) was added 1N NaOH (80 ml, 80 mmol). After stirring for 1 h, the solvents were evaporated and the residue was dissolved in 1N HCl (80 ml, 80 mmol) and then concentrated, azeotroped with CH₃CN to give the crude acid. The crude acid was suspended in DMF (200 mL) and then treated with HCl•HN(Me)OMe (13.9 g, 144 mmol), EDC (15.1g, 79.2 mmol), HOBT (9.6g, 72 mmol) and NMM (60 mL, 576 mmol). The mixture was stirred for 18 hours and then concentrated. The residue was dissolved in ethyl acetate, washed with 10% K₂CO₃, brine, dried (MgSO₄), and concentrated to give amide 15-3 as a brown oil.

TLC R_f = 0.30 (silica, 70:25:5 chloroform/ ethyl acetate/ MeOH)

5-Ethoxy-pyridine-3-carbaldehyde (15-4)

To a stirred solution of 15-3 (14.0 g, 66.5 mmol) and CH₂Cl₂ (200 mL) at -78°C under argon was added DIBAL (1.0M hexanes, 90ml) dropwise over 30 minutes. After 30 minutes, the solution was warmed to 0°C for 1 hour. The reaction was quenched with 100 ml 1.0M Rochelle's salt, stirred for 1.0 hour and then extracted with Et₂O. The organic layer was dried (MgSO₄), and then concentrated to give the aldehyde 15-4 as a brown oil.

TLC R_f = 0.32 (silica, 70:25:5 chloroform/ethyl acetate/MeOH)

¹H NMR (300 MHz, CDCl₃) δ 10.10 (s, 1H), 8.65 (s, 1H), 8.55 (s, 1H), 7.59 (s, 1H), 4.14 (q, 2H, J=7Hz), 1.43 (t, 3H, J=7Hz).

3-(5-Ethoxy-pyridin-3-yl)-acrylic acid *tert*-butyl ester (15-6)

A mixture of 15-4 (8.0 g, 51.6 mmol), 15-5 (20 g, 54.2 mmol), and benzene (150 mL) was heated to reflux for 30 minutes. The mixture was diluted with Et₂O and then washed with 10% K₂CO₃, brine and dried (MgSO₄). Following evaporative removal of the solvent, the residue was chromatographed (silica gel, 30% EtOAc/hexanes) to give 15-6 as a yellow solid.

TLC R_f = 0.41 (silica, 70:25:5 chloroform/ ethyl acetate/ MeOH)

¹H NMR (300 MHz, CDCl₃) δ 8.31 (m, 2H), 7.55 (d, 1H, J=16Hz), 7.27 (s, 1H), 6.40 (d, 1H, J=16Hz), 4.10 (q, 2H, J=7Hz), 1.54 (s, 9H), 1.44 (m, 3H).

3(S)-Amino-3-(5-ethoxy-pyridin-3-yl)-propionic acid *tert*-butyl ester (15-8)

To a stirred solution of 15-7 (500mg, 2.38mmol) and THF at 0°C was added nBuLi (2.5 M THF, 0.95 ml) dropwise. After 20 minutes, the solution was cooled to -78°C and 15-6 (500mg, 1.98 mmol), dissolved in 3 ml THF, was added. After 15 minutes, the reaction was quenched with sat. NH₄Cl followed by the removal of the cooling bath. The solution was extracted with ethyl acetate. The organic portion was washed with brine, dried (MgSO₄) and concentrated. The residue was dissolved in acetic acid (14 ml), and the solution was purged with argon for 30 minutes. 10% Pd/C (1.0 g) was added and the mixture was heated to 80°C. 1,4-Cyclohexadiene (6 ml) was added dropwise maintaining an internal temperature between 80°C and 90°C. After 5.0 hours, the mixture was filtered through a celite pad, concentrated and then azeotroped with toluene. The residue was chromatographed (silica gel, 5% [10:10:1 EtOH/NH₄OH/H₂O]/ 70:25:5 chloroform/ ethyl acetate/ MeOH) to give 15-8 as a yellow solid.

¹H NMR (300 MHz, CDCl₃) δ 8.18 (m, 2H), 7.25 (s, 1H), 4.41 (m, 1H), 4.08 (q, 2H, J=7Hz), 2.59 (m, 2H), 1.87 (s, 2H), 1.40 (m, 12H).


3(S)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid (15-9)

The title compound 15-9 was prepared from 2-9A and 15-8 using the procedure described in Scheme 2.

TLC R_f = 0.27 (silica, 10:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

^1H NMR (300 MHz, CD_3OD) δ 8.13 (m, 2H), 7.48 (d, 1H, $J=7\text{Hz}$), 7.35 (s, 1H), 6.55 (d, $J=8\text{Hz}$, 1H), 5.53 (m, 1H), 4.13 (q, 2H, 7Hz), 3.31-3.70 (m, 7H), 3.06 (m, 2H), 2.55-2.85 (m, 6H), 1.88-2.15 (m, 5H), 1.42 (t, 3H, $J=7\text{Hz}$).

SCHEME 16



3(S)-Amino-3-(5-methoxy-pyridin-3-yl)-propionic acid tert-butyl ester (16-2)

3-Bromo-5-methoxy-pyridine 16-1 (prepared as described in *J. Org. Chem.* 1990, 55, 69) was converted into 16-2 utilizing the procedure previously described for the conversion of 1-4 to 1-6.
5 ¹H NMR (300 MHz, CD₃OD) δ 8.20 (d, 1H, J=3Hz), 8.18 (d, 1H, J=2Hz), 7.50 (s, 1H), 4.51 (m, 1H), 3.90 (s, 3H), 2.87 (m, 2H), 1.37 (m, 9H).

10 3(S)-(5-Methoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl ester (16-3)

The title compound 16-3 was prepared from 2-9A and 16-2 using the procedure described in Scheme 2.
15 TLC R_f = 0.27 (silica, 70:20:10 chloroform/ethyl acetate/MeOH)
¹H NMR (300 MHz, CDCl₃) δ 8.23 (d, 1H, J=3Hz), 8.15 (s, 1H), 7.22 (s, 1H), 7.02 (d, 1H, J=7Hz), 6.33 (d, 1H, 7Hz), 5.46 (t, 1H, J=8Hz), 4.78 (s, 1H), 4.11 (m, 2H), 3.84 (s, 3H), 3.30 (m, 6H), 3.00 (m, 2H), 2.67 (t, 2H, J=6Hz), 2.52 (m, 2H), 1.85 (m, 6H), 1.23 (m, 3H).

20 3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid ethyl ester (16-4)

To a stirred solution of 16-3 (200 mg, 0.4278mmol) and ethanethiol (0.5ml) and CH₂Cl₂ (3 ml) was added AlCl₃ (570mg, 4.28 mmol). After 1.0 hour, the reaction was quenched with sat. NaHCO₃. Ethyl acetate was added, and the reaction mixture was then purged with argon for 1.0 hour. The organic layer was separated, washed with brine, dried (MgSO₄) and then concentrated to give the phenol 16-4 as a
25 yellow oil.
30 TLC R_f = 0.22 (silica, 70:20:10 chloroform/ethyl acetate/MeOH)


3(S)-(5-Hydroxy-pyridin-3-yl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (16-5)

The title compound 16-5 was prepared from 16-4 by basic hydrolysis using the procedure described in Scheme 2.

5 TLC Rf = 0.39 (silica, 10:1:1 EtOH/water/NH₄OH).

¹H NMR (300 MHz, CD₃OD) δ 8.01 (m, 2H), 7.46 (d, 1H, J=7Hz), 7.20 (s, 1H), 6.53 (d, J=8 Hz, 1H), 5.49 (m, 1H), 3.51-3.68 (m, 2H), 3.46 (t, 2H, 5Hz), 3.19 (m, 2H), 3.00 (m, 2H), 2.52-2.78 (m, 6H), 1.92 (m, 4H).

SCHEME 17



3(S)-(Ethynyl)-3-[2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-propionic acid (17-2)


15 The title compound 17-2 was prepared from 2-9A and 17-1 (for preparation see J.A. Zablocki, et.al., *J. Med. Chem.* 1995, 38, 2378-2394) using the procedure described in Scheme 2.

TLC Rf = 0.32 (silica, 15:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

¹H NMR (300 MHz, CD₃OD) δ 7.45 (d, J=7Hz, 1H), 6.53 (d, J=8 Hz, 1H), 5.15 (m, 1H), 3.31-3.70 (m, 7H), 2.55-2.85 (m, 7H), 2.35 (m, 1H), 1.88 -2.15 (m, 4H).

20

SCHEME 18




2(S)-Benzenesulfonylamino-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid (18-2)

The title compound 18-2 was prepared from 2-9A and 18-1
 (for preparation, see Scheme A substituting phenylsulfonyl chloride for
 5 4-iodophenylsulfonyl chloride) using the procedure described in Scheme
 2.

TLC R_f = 0.23 (silica, 15:10:1:1 ethyl acetate/EtOH/water/NH₄OH).

¹H NMR (300 MHz, CD₃OD) δ 7.81 (m, 2H), 7.36 (m, 3H), 7.10 (d, 1H,
 J=8Hz), 6.37 (d, 1H, J=7Hz), 3.61 (m, 1H), 3.36 (m, 2H), 3.02 & 3.18 (m,
 10 6H), 3.00 (m, 2H), 2.68 (t, 2H, J=6Hz), 2.50 (m, 2H), 1.79 & 1.90 (m, 4H).

SCHEME 19



3-Amino-pent-4-enoic acid ethyl ester (19-1)

15 A mixture of 5% Pd/BaSO₄ (0.025 g) and quinoline (0.30 mL)
 was stirred under a balloon of hydrogen for 30 minutes. 3-Amino-pent-4-
 ynoic acid ethyl ester 17-1 (1.77 g, 10.0 mmol) in EtOH (15 mL) was added
 and the solution stirred for an additional 2.5 hours. The solution was
 filtered through a pad of celite and concentrated in vacuo to provide 2.65
 20 g of crude product 19-1.

¹H NMR (CDCl₃, 300 MHz): δ 8.40-7.60 (br s, 2H), 6.11-5.96 (m, 1H), 5.58-5.53 (d, 1H), 5.44-5.41 (d, 1H), 4.31-4.16 (m, 3H), 3.12-2.86 (m, 2 H), 1.29-1.25 t, 3H).

5 3(S)-[2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl]-pent-4-enoic acid (19-2)

The title compound 19-2 was prepared from 2-9A and 19-1 using the procedure described in Scheme 2.

10 ¹H NMR (CDCl₃, 300 MHz): δ 11.1 (s, 1H), 7.21-7.19 (d, 1H), 6.26-6.23 (d, 1H), 5.91-5.78 (m, 1H), 5.22-5.00 (m, 3H), 3.79-3.16 (m, 10H), 2.77-2.33 (m, 5H), 2.06-1.80 (m, 4H).

MS (FAB) 359 (M+1)

20-1 $\xrightarrow[\text{CuI, NEt}_3]{\text{Pd(PPh}_3)_4}$ 20-2 $\xrightarrow[\text{H}_2, \text{EtOH}]{\text{Et}_3\text{N, PtO}_2}$ 20-3, R = BOC

20-3, R = BOC $\xrightarrow[\text{NaCNBH}_3]{\text{DIPEA, AcOH}}$ 20-4, R = NH₂·HCl

20-4, R = NH₂·HCl $\xrightarrow[\text{THF}]{\text{BOC}_2\text{O}}$ 20-5, R = H

20-5, R = H $\xrightarrow[\text{THF}]{\text{BOC}_2\text{O}}$ 20-6, R = BOC


20-6, R = BOC $\xrightarrow[\text{MeNHOMe}\cdot\text{HCl}]{\text{NaOH, EDC, HOBT, MM}}$ 20-7

20-7 $\xrightarrow[\text{THF}]{\text{DIBAL}}$ 20-8

20-8 $\xrightarrow[\text{isopropanol}]{\text{15-8, AcOH, NaOAc, NaCNBH}_3}$ 20-9, R = BOC

20-9, R = BOC $\xrightarrow[\text{CH}_2\text{Cl}_2]{\text{p-TSA}}$ 20-10, R = H

SCHEME 20 (CONTINUED)



[3-(6-Bromo-pyridin-2-yl)-propyl]-(2-oxo-ethyl)-carbamic acid *tert*-butyl ester (20-2)

A solution of 2,6-dibromopyridine 20-1 (111 g, 468 mmol) and N-BOC-propargylamine (80.0 g, 515 mmol) in 500 ml of triethylamine at 0°C was treated with copper(I) iodide (2.23 g, 11.7 mmol). The mixture was purged with argon and then dichlorobis(triphenylphosphine)-palladium(II) (8.22 g, 11.7 mmol) was added. The solution was stirred at 0°C for one hour, then at room temperature for 16 hours. The solution was diluted with 250 mL ether and washed with H₂O (4 X 100 mL). The organic extract was washed with brine and dried over Na₂SO₄. The solvents were removed in vacuo and the crude product was purified by silica gel chromatography (25% EtOAc/hexane) to afford 20-2.
1H NMR (CDCl₃, 300 MHz): δ 7.53-7.34 (m, 3 H), 4.82-4.80 (br s, 1 H), 4.18-4.17 (d, 2 H), 1.46 s, 9 H).

3-(2-([3-(6-Bromo-pyridin-2-yl)-propyl]-*tert*-butoxycarbonyl-amino)-ethylamino) (20-3)

To a solution of 20-2 (79.8 g, 257 mmol) in 350 mL of ethanol and triethylamine (26.8 mL, 193 mmol) was added platinum(IV) oxide (2.91 g, 12.8 mmol). After stirring under a hydrogen atmosphere for 4 hours, the solution was filtered through a pad of celite and concentrated in vacuo. The crude product was dissolved in EtOAc (200 mL) and washed with H₂O (4 X 250 mL) and brine (250 mL), dried over Na₂SO₄, filtered, and concentrated in vacuo. The crude material was purified by silica gel chromatography (10% EtOAc/CHCl₃) to afford 20-3.
1H NMR (CDCl₃, 400 MHz): δ 7.48-7.42 (t, 1 H), 7.32-7.29 (d, 1H), 7.13-7.10 (d, 1H), 4.71-4.70 (br s, 1 H), 3.18-3.09 (m, 2 H), 2.82-2.77 (t, 2 H), 1.96-1.85(m, 2 H), 1.44 (s, 9 H).

3-(6-Bromo-pyridin-2-yl)-propylamine•hydrochloride (20-4)

A solution of 20-3 (3.33 g, 10.5 mmol) in EtOAc (150 mL) was saturated with HCl gas and stirred at room temperature for 2 hours. The solvent was removed in vacuo to provide 20-4. The crude product was used in the next step without further purification.

[3-(6-Bromo-pyridin-2-yl)-propylamino]-acetic acid ethyl ester (20-5)

A solution of 3-(6-bromo-pyridin-2-yl)-propylamine 20-4 (25.6 g, 89.1 mmol), diisopropylethylamine (46.5 mL, 267 mmol), acetic acid (28 mL, 490 mmol), and ethyl glyoxylate (10.9 g, 107 mmol) in 200 mL of methanol was stirred at room temperature for one hour. A 1M solution of NaCNBH₃ in THF (98.0 mL, 98.0 mmol) was added slowly with a syringe pump over 4 hours. The resulting solution was stirred for 12 hours, after which the solvent was removed in vacuo and the residue taken up in chloroform and filtered. The solution was then washed with 10% Na₂CO₃, dried over Na₂SO₄, and the solvent removed in vacuo to give the crude amine. The crude product was purified by silica gel chromatography (7% MeOH/CHCl₃) to give 20-5 in a 3:2 mixture of ethyl and methyl esters.

¹H NMR (CDCl₃, 300 MHz) δ 7.47-7.42 (t, 1H), 7.31-7.27, (t, 1H), 7.13-7.10 (d, 1H), 4.20-4.14, (m, 2H), 3.39 (s, 2H), 2.85-2.75 (m, 2H), 2.68-2.63 (t, 2H), 1.96-1.88 (m, 2H), 1.29-1.24 (m, 3H).

[[3-(6-Bromo-pyridin-2-yl)-propyl]-tert-butoxycarbonyl-amino]-acetic acid ethyl ester (20-6)

To a solution of [3-(6-bromo-pyridin-2-yl)-propylamino]-acetic acid ethyl ester 20-5 (17.6 g, 58.6 mmol) in THF (200 mL) was added di-*tert*butyldicarbonate (15.3 g, 70.3 mmol). After stirring at room temperature for 16 hours, the solvents were removed in vacuo. The product was purified by silica gel chromatography (5% MeOH/CHCl₃) to give 20-6.

¹H NMR (CDCl₃, 300 MHz) δ 7.47-7.42 (m, 1H), 7.32-7.28 (t, 1H), 7.16-7.10 (t, 1H), 4.22-4.15, (q, 2H), 3.95-3.85 (d, 2H), 3.38-3.29 (m, 2H), 2.80-2.75 (t, 2H), 2.03-1.91 (m, 2H), 1.46-1.44 (m, 9H), 1.31-1.23 (m, 3H).

[3-(6-Bromo-pyridin-2-yl)-propyl]-[(methoxy-methyl-carbamoyl)-methyl]-carbamic acid *tert*-butyl ester (20-7)

To a solution of 20-6 (23.4 g, 58.4 mmol) in ethanol (200 mL) was added NaOH (100 mL 1M solution in water, 100 mmol). After stirring for 1 h at 50°C, HCl (10.3 mL 12 M, 4.75 mmol) was added in 50 mL EtOH, and the mixture evaporated to give an oily residue. The

residue was evaporated from ethanol three times, and then from acetonitrile three times, producing a yellow crusty solid which was dried under a vacuum of <2 mm Hg for 2 h. This residue was then slurried in acetonitrile (180 mL) and chloroform (180 mL), and NMM (41.7 mL, 379
5 mmol), N,O-dimethylhydroxylamine hydrochloride (11.9 g, 122 mmol), HOBT (10.2 g, 75.9 mmol), and EDC (14.5 g, 75.9 mmol) were added. After stirring for 15 h, the mixture was evaporated to dryness, the residue slurried in EtOAc, washed with sat. NaHCO₃, brine, and dried over Na₂SO₄. Evaporative removal of the solvent followed by evaporation
10 from toluene to remove the residual NMM gave 20-7 as a yellow oil. TLC R_f = 0.49 (silica, 70:25:5 chloroform/ethyl acetate/methanol) ¹H NMR (CDCl₃, 300 MHz) δ 7.48-7.41 (m, 1H), 7.32-7.28 (t, 1H), 7.17-7.11 (t, 1H), 4.14 (s, 2H), 3.73-3.70 (d, 3H), 3.39-3.30 (m, 2H), 3.18 (s, 3H), 2.80-2.75 (t, 2H), 2.02-1.91 (m, 2H), 1.45 (m, 9H).

15

[3-(6-Bromo-pyridin-2-yl)-propyl]-(2-oxo-ethyl)-carbamic acid tert-butyl ester (20-8)

To a stirred solution of 20-7 (14.9 g, 35.7 mmol) and THF (100 ml) at -78°C was added DIBAL (1.0M/hexanes, 53.6 ml, 53.6 mmol)
20 dropwise over 20 minutes. After 1 h, the mixture was warmed to RT and quenched by the careful addition of 20 mL MeOH. 200 ml of 1.0 M Rochelle's salt was then added followed by the removal of the cooling bath. The mixture was stirred for 1.0 hour and then diluted with Et₂O. After another 30 minutes of stirring, the organic portion was separated
25 and dried over MgSO₄. Evaporative removal of the solvent gave the crude aldehyde 20-8 as a colorless oil. ¹H NMR (CDCl₃, 300 MHz) δ 9.59-9.56 (d, 1H), 7.48-7.43 (t, 1H), 7.32-7.26 (m, 1H), 7.14-7.07 (m, 1H), 3.53-3.26 (m, 4H), 2.80-2.72 (m, 2H), 2.00-1.93 (m, 2H), 1.44 (s, 9H).

30

3-(5-Ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester (20-9)

A mixture of the crude aldehyde 20-8 (0.671 g, 1.88 mmol), amine 15-8 (0.651 g, 2.44 mmol), acetic acid (0.107 mL, 1.88 mmol), NaOAc (1.54 g, 18.8 mmol), and powdered molecular sieves (1.20g) in 2-propanol (15 mL) was stirred for 20 minutes. The mixture was cooled to
35

0°C and then NaBH₃CN (0.354 g, 5.64 mmol) was added. After stirring 6 hours, the pH of the mixture was adjusted to ~2 with 1N HCl. The solution was stirred for an additional 10 minutes, ethyl acetate (20 mL) was added, and the pH was adjusted to ~11 with 10% potassium carbonate. The organics were extracted with ethyl acetate, dried over Na₂SO₄, and removed in vacuo. The residue was chromatographed (silica gel, [70:25:5 CHCl₃/EtOAc/MeOH]) to give 20-9 as a yellow oil in 95% yield.

¹H NMR (CDCl₃, 300 MHz): δ 8.19-8.18 (d, 1H), 8.14-8.13 (d, 1H), 7.48-7.42 (t, 1H), 7.32-7.26 (m, 1H), 7.23-7.20 (m, 1H), 7.11-7.08 (m, 1H), 4.16-4.02 (m, 3H), 3.30-3.18 (m, 4H), 2.74-2.45 (m, 7H), 1.92-1.86 (t, 2H), 1.45-1.38 (m, 21H).

15 3-[2-[3-(6-Bromo-pyridin-2-yl)-propylamino]-ethylamino]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester (20-10)

To a solution of 3-(5-ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester 20-9 (0.085g, 0.141 mmol) in dichloromethane (5 mL) was added *p*-toluenesulfonic acid (0.161 g, 0.847 mmol). The mixture was stirred for 2 hours at room temperature and was then neutralized with 1N NaOH. The organic layer was extracted (3 X 25 mL) with CHCl₃, dried, and concentrated in vacuo. The crude product 20-10 was not purified (0.069 g, 96% yield).

¹H NMR (CDCl₃, 300 MHz): δ 8.23-8.21 (m, 2H), 7.45 (s, 1H), 7.40-7.35 (m, 1H), 7.28-7.25 (t, 1H), 7.04-7.01 (d, 1H), 6.28 (br s, 2H) 4.39-4.33 (t, 1H), 4.00-3.92 (q, 2H), 3.40-3.35 (m, 2H), 3.28-3.22 (m, 1H), 3.15-2.90 (m, 4H), 2.79-2.71 (m, 3H), 2.14-2.01 (m, 2H), 1.34-1.26 (m, 12H).

3-[3-[3-(6-Bromo-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid tert-butyl ester (20-11)

30 . To a stirred solution of 20-10 (0.80 g, 1.57 mmol) and diisopropylethylamine (0.823 mL, 4.72 mmol) in dichloromethane (10 mL) at 0°C was added *p*-nitrophenyl chloroformate (0.333 g, 1.65 mmol). The solution stirred for 30 minutes and dioxane (10 mL) was added, then refluxed for 4 hours. EtOAc (100 mL) was added and the organics were

washed with 10% K₂CO₃, dried, and concentrated in vacuo. The residue was chromatographed (silica gel, [70:20:10 CHCl₃/ EtOAc/ MeOH]) to give 20-11.

¹H NMR (CDCl₃, 400 MHz) δ 8.22-8.18 (dd, 1H), 8.14-8.13 (t, 1H), 7.46-7.37 (m, 1H), 7.31-7.24 (m, 1H), 7.20-7.16 (m, 1H), 7.11-7.09 (d, 1H), 4.09-4.04 (q, 2H), 3.34-3.16 (m, 5H), 2.99-2.87 (m, 2H), 2.77-2.69 (m, 2H), 2.63-2.46 (m, 2H), 1.97-1.88 (m, 2H), 1.44-1.37 (m, 12H).
MS M+1 = 533.3

10 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-(3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl)-2-oxo-imidazolidin-1-yl)-propionic acid *tert*-butyl ester (20-12)

To a stirred solution of 20-11 (0.075 g, 0.142 mmol) in toluene (3 mL) was added Pd(DBA)₂ (0.0041 g, 0.0071 mmol), DPPF (0.0039 g, 0.0071 mmol), and NaOt-Bu (0.0163 g, 0.170 mmol) followed by *p*-methoxybenzylamine (0.0204 mL, 0.156 mmol). The resulting solution was heated at 110°C for 2 hours. The solution was cooled and the solvent was removed in vacuo. The product was purified by silica gel chromatography (10% EtOH/EtOAc) to give 20-12.

¹H NMR (CDCl₃, 400 MHz) δ 8.21-8.20 (d, 1H), 8.14-8.13 (d, 1H), 7.33-7.25 (m, 3H), 7.19-7.17 (t, 1H), 6.87-6.84 (d, 2H), 6.45-6.43 (d, 1H), 6.21-6.18 (d, 1H), 5.48-5.42 (t, 1H), 5.26 (br s, 1H), 4.38-4.37 (d, 2H), 4.09-4.01 (q, 2H), 3.79 (s, 3H), 3.31-3.18 (m, 5H), 3.07-2.87 (m, 3H), 2.63-2.58 (t, 2H), 1.95-1.84 (m, 2H), 1.49-1.39 (m, 12H).

25 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-(3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl)-2-oxo-imidazolidin-1-yl)-propionic acid (20-13)

To a stirred solution of 20-12 (0.028 g, 0.047 mmol) in dichloromethane (10 mL) was added TFA (1 mL). After 1 hour, the solvent was removed in vacuo and azeotroped twice with toluene (15 mL). The residue was chromatographed (silica gel, 25:10:1:1 followed by 15:10:1:1 ethyl acetate /EtOH /water/NH₄OH) to give 20-13 as a white solid.

¹H NMR (CDCl₃, 400 MHz): δ 8.21-8.20 (d, 2H), 7.50-7.46 (t, 1H), 7.31-7.26 (m, 2H), 7.19 (s, 1H), 6.88-6.85 (d, 2H), 6.42-6.40 (d, 1H), 6.34-6.32 (d, 1H),

5.69-5.63 (m, 1H), 5.30 (s, 1H), 4.43-4.41 (d, 2H), 4.09-4.01 (q, 2H), 3.77 (s, 3H), 3.75-3.44 (m, 3H), 3.24-2.86 (m, 4H), 2.79-2.67 (m, 3H), 2.03-1.90 (m, 2H), 1.44-1.41 (t, 3H).

MS (FAB) 534 (M+1)

5


3-{3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl}-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid (20-14)

To a stirred solution of 20-13 (0.031 g, 0.052 mmol) in dichloromethane (10 mL) was added TFA (1 mL). The solution was stirred for 16 hours at 85°C. after which the solvent was removed in vacuo and azeotroped twice with toluene (15 mL). The residue was chromatographed (silica gel, 15:10:1:1 followed by 10:10:1:1 ethyl acetate /EtOH /water /NH₄OH) to give 20-14 as a white solid.


¹H NMR (CD₃OD, 400 MHz): δ 8.13-8.10 (m, 2H), 7.59-7.54 (t, 1H), 7.38-7.35 (m, 1H), 6.60-6.57 (d, 2H), 5.53-5.47 (q, 1H), 4.15-4.09 (q, 2H), 3.64-3.57 (m, 1H), 3.47-3.41 (m, 1H), 3.28-3.21 (m, 2H), 3.07-2.90 (m, 3H), 2.76-2.61 (m, 3H), 2.02-1.83 (m, 2H), 1.42-1.38 (t, 3H).

MS (FAB) 414 (M+1)

Scheme 21



SCHEME 21 (CONTINUED)

3-Bromo-6-chloro-5-nitropyridine (21-2)

A suspension of CuCl_2 (3.33 g, 24.8 mmol) in anhydrous
 5 CH_3CN (200 mL) at 65° was treated with tert-butyl nitrite (3.13 mL, 26.3
 mmol), followed by the dropwise addition of a solution of 21-1 in 60 mL of
 CH_3CN . The resulting mixture was stirred under an argon
 atmosphere at 65° for 2 h and concentrated at reduced pressure. The
 residue was partitioned between EtOAc (150 mL) and 3% HCl (60 mL),
 10 and the organic layer washed successively with 3% HCl, water, and
 brine (60 mL), then dried, filtered and concentrated to afford a brown
 solid which was chromatographed on silica (25% EtOAc/Hexane) to
 afford 21-2 as a yellow crystalline solid.

TLC R_f = 0.60 (25% EtOAc/Hexane)

15 ^1H NMR (300 MHz, CDCl_3) δ 8.70 (d, J =2.4 Hz, 1H), 8.37 (d, J =2.4 Hz, 1H).

(3-Nitro-5-bromo-pyridin-2-yloxy)-acetic acid methyl ester (21-3)

Methyl glycolate (450 mg, 5.05 mmol) was added to a suspension of 60% NaH (131 mg, 55 mmol) in THF (20 mL) at 0°. The resulting solution was stirred under argon for 0.5 h, then treated with a solution of 21-2. After stirring at 0° for 0.5 h, the reaction was diluted with ethyl acetate, and washed successively with sat. NaHCO₃, water and brine (80 mL each), then dried, filtered and concentrated to afford 21-3 as a yellow solid.

TLC Rf = 0.70 (25% EtOAc/ Hexane)

¹H NMR (300 MHz, CDCl₃) δ 8.46 (d, J= 2.4 Hz, 1H), 8.37 (d, J= 2.4 Hz, 1H) 5.15 (s, 2H), 3.78 (s, 3H).

2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-7-bromo-naphthalene (21-4)

A mixture of 21-3 (1.5 g, 5.12 mmol) and powdered tin (1.37 g, 11.5 mmol) was treated with conc. HCl (10 mL). The mixture was heated to 80° for 2 h, then cooled and concentrated. The residue was partitioned between CHCl₃ and sat. NaHCO₃, washed with brine, then dried, filtered and concentrated to afford a yellow solid.

Chromatography on silica gel (50% hexane/EtOAc) gave 21-4 as a yellow solid.

TLC Rf = 0.65 (50% EtOAc/ Hexane)

¹H NMR (300 MHz, DMSO-d₆) δ 10.81 (br,s, 1H), 7.88 (d, J=2.4 Hz, 1H), 7.25 (d, J=2.4 Hz, 1H), 4.81 (s, 2H).

3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-acrylic acid tert-butyl ester (21-5)

A mixture of 21-4 (1.12 g, 4.89 mmol), (o-tol)₃P (298 mg, 1.0 mmol), Pd(OAc)₂ (110 mg, 0.49 mmol), and triethylamine (0.86 mL, 5.87 mmol) in DMF (20 mL) was placed in a 100-mL flask. The mixture was degassed with argon, then tert-butyl acrylate (752 mg, 5.87 mmol) was added and the tube sealed and heated to 100° for 12 h. The reaction mixture was diluted with ethyl acetate, filtered and washed with NaHCO₃, water, and brine, dried, filtered and concentrated.

Chromatography on silica gel (25% hex/EtOAc) gave 21-5 as a yellow solid.

TLC Rf = 0.60 (25% EtOAc/ Hexane)

¹H NMR (300 MHz, DMSO-d₆) δ 10.91 (br,s, 1H), 8.15 (d, J=2.4 Hz, 1H), 7.54 (d, J=16 Hz, 1 H), 7.42 (d, J= 2.4 Hz, 1H), 6.35 (d, J=16 Hz, 1 H), 4.84 (s, 2H), 1.48 (s, 9H).

5 3(S)-[Benzyl-(1(R)-phenylethyl)-aminol]-3-(2-oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert*-butyl ester (21-6)

A solution of N-benzyl-α-(R)-methylbenzylamine (0.82 g, 3.87 mmol) in THF (25 mL) at 0°C was treated with n-BuLi (1.6 mL of a 2.5 M soln in hexanes). The resulting solution was stirred at 0°C for 30 min and then cooled to -78°C. A solution of acrylate 21-5 (0.485 g, 1.76 mmol) in THF (5 mL) was added. After stirring for 15 min at -78°C, satd aq NH₄Cl soln (5 mL) was added and the cold bath removed. The mixture was warmed to room temperature, and extracted with Et₂O (2 x 40 mL). The combined organic extracts were washed with brine (30 mL), dried over MgSO₄, filtered, and concentrated. The residue was purified by flash chromatography (40% ethyl acetate/hexanes) to give the β-aminoester 21-6 as a yellow oil.

TLC R_f = 0.3 (40% ethyl acetate/hexanes)

¹H NMR (300 MHz, CDCl₃) δ 1H NMR 8.70 (br, s, 1H), 7.91 (d, J=1.8 Hz, 1H), 7.4-7.2 (10H), 7.12 (d, J=1.8 Hz, 1H), 4.80 (s, 2 H), 4.42 (m, 1H), 3.91 (q, J=6.7 Hz, 1 H), 3.69 (d, J=7.2 Hz, 1H,), 3.62 (d, J =7.2 Hz, 1H,), 2.46 (m, 2H), 1.34 (d, J=7.0 Hz, 3H), 1.29 (s, 9H).

25 3(S)-Amino-3-(2-oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert*-butyl ester (21-7)

A mixture of the dibenzylamine 21-6 (0.22 g, 0.44 mmol) in EtOH/H₂O/AcOH (26 mL/3 mL/1.0 mL) was degassed with argon and treated with Pd(OH)₂ (100 mg). The mixture was placed under 1 atm of H₂. After stirring for 18 h, the mixture was diluted with EtOAc and filtered through celite. The filtrate was concentrated and the residue purified by flash chromatography (20% 20:1:1 EtOH/NH₄OH/H₂O - 80% EtOAc) to give the *tert*-butyl ester 21-7 as a white solid.

TLC R_f = 0.5 (20% 20:1:1 EtOH/NH₄OH/H₂O - 80% EtOAc)

¹H NMR (300 MHz, CD₃OD) δ 7.89 (d, J= 1.7 Hz, 1H), 7.31 (d, J=1.7 Hz, 1H), 4.81 (s, 2H), 4.38 (m, 1H), 2.6, (m, 2H), 1.41 (s, 9H).

5 3(R)-[Benzyl-(1-phenylethyl)-aminol-3(S)-(2-thioxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert.* butyl ester (21-8)

A solution of 21-6 (0.22 g, 0.44 mmol) in anhydrous THF was treated with Lawesson's reagent (0.098 g, 0.243 mmol) and stirred at room temperature for 1.5 h. Silica gel (500 mg) was added to the reaction mixture and the solvent was removed at reduced pressure and the product was eluted from silica using 25% EtOAc/hexane to afford 21-8 as a yellow solid.

TLC R_f = 0.7 (40% EtOAc/hexane)

¹H NMR (300 MHz, CD₃OD) δ 9.82 (br, s, 1H), 7.95 (d, J=1.8 Hz, 1H), 7.4-7.2 (11H), 5.08 (s, 2 H), 4.42 (m, 1H), 3.91 (q, J=6.7 Hz, 1 H), 3.69(d, J=7.2 Hz, 1H,), 3.62 (d, J =7.2 Hz, 1H), 2.46 (m, 2H), 1.34 (d, J=7.0 Hz, 3H), 1.29 (s, 9H).

3(S)-Amino-3-(2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-propionic acid *tert*-butyl ester (21-9)

20 A solution of 21-8 (1.0 g, 1.9 mmol) in anhydrous Et₂O (10 mL) at 0° was treated dropwise with LiAlH₄ (2.09 ml of a 1.0 M solution in Et₂O). The resulting solution was stirred at 0°C for 30 min and then quenched by the sequential addition of H₂O (0.3 mL), 15 % NaOH (0.08 mL). Celite (1 g) was added and the mixture filtered through a Celite pad. The filtrate was evaporated and the residue was purified by flash chromatography (65% ethyl acetate/hexanes) to give the dibenzylamine intermediate as a yellow oil.

TLC R_f = 0.4 (65% ethyl acetate/hexanes)

¹H NMR (300 MHz, CDCl₃) δ ¹H NMR 7.61 (d, J=1.8 Hz, 1H), 7.4-7.2 (10H), 6.87 (d, J=1.8 Hz, 1H), 4.41 (m, 2 H), 4.36 (m, 1H), 3.91 (q, J=6.7 Hz, 1 H), 3.8 (brs, 1H), 3.69 (m, 2H), 3.42 (m, 2H), 2.46 (m, 2H), 1.34 (d, J=7.0 Hz, 3H), 1.29 (s, 9H).

This material was deprotected with Pd(OH)₂ in ethanol to afford 21-9 as a white solid. TLC R_f = 0.5 (20% 20:1:1 EtOH/NH₄OH/H₂O - 80% EtOAc)

¹H NMR (300 MHz, CD₃OD) δ 7.59 (d, J=1.7 Hz, 1H), 6.92 (d, J=1.7 Hz, 1H), 4.41 (m, 2H), 4.30 (m, 1H),), 3.41 (m, 2H), 2.6, (m, 2H), 1.41 (s, 9H).

5 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (21-10)

The title compound 21-10 was prepared from 2-9A and 21-7 using the procedure described in Scheme 2.

High resolution MS Calc'd.=418.2198, Obs'd = 481.2193.


10

3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (21-11)

15 The title compound 21-11 was prepared from 2-9A and 21-8 using the procedure described in Scheme 2.

High resolution MS Calc'd.=467.2417, Obs'd = 467.2401.

Scheme 22



3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-7-bromo-naphthalene (22-2)

A solution of 22-1 (4.8 g, 32 mmol) in MeOH (160 mL) at -15° was treated dropwise with bromine (25.7 g, 161 mmol). After stirring at -15° for 0.5 h, the mixture was warmed to ambient temperature and stirred overnight. The resulting white precipitate was filtered and washed with cold MeOH to afford 22-2 as a white solid.

TLC R_f = 0.65 (50% EtOAc /Hexane)

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 11.2 (br s, 1H), 8.05 (d, J = 2.4 Hz, 1H), 7.66

(d, J = 2.4 Hz, 1H), 4.76 (s, 2H).

3(S)-Amino-3-(3-oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-propionic acid tert-butyl ester (22-3)

Bromide 22-2 was converted to amino ester 22-3 as

5. illustrated in Scheme 21.

TLC R_f = 0.5 (12% 20:1:1 EtOH/NH₄OH/H₂O - 88% EtOAc)

¹H NMR (300 MHz, CD₃OD) δ 8.04 (d, J = 1.7 Hz, 1H), 7.34 (d, J = 1.7 Hz, 1H), 4.76 (s, 2H), 4.38 (m, 1H), 2.6, (m, 2H), 1.41 (s, 9H).

10 3(S)-Amino-3-(3-oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-propionic acid tert-butyl ester (22-4)

Bromide 22-2 was converted to amino ester 22-4 as illustrated in Scheme 21.

TLC R_f = 0.5 (20% 20:1:1 EtOH/NH₄OH/H₂O - 80% EtOAc)

15 ¹H NMR (300 MHz, CD₃OD) δ 8.04 (d, J = 1.7 Hz, 1H), 7.34 (d, J = 1.7 Hz, 1H), 4.76 (s, 2H), 4.38 (m, 1H), 2.6, (m, 2H), 1.41 (s, 9H).

3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid (22-5)

20

The title compound 22-5 was prepared from 2-9A and 22-3 using the procedure described in Scheme 2.


High resolution MS Calc'd.=481.2198, Obs'd = 481.2194.

25 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid (22-6)

The title compound 22-6 was prepared from 2-9A and 22-4 using the procedure described in Scheme 2.

30 High resolution MS Calc'd.=467.2417, Obs'd = 467.2411.

Scheme 23

Furo-[2,3-b]pyridine-5-carboxaldehyde (23-2)

- A solution of alcohol **23-1** (M. Bhupathy, et al., J. Heterocycl. Chem. 1995, 32, 1283-1287) was treated with excess MnO_2 (10 eq) and the mixture stirred at room temperature for 16 h, then filtered through Celite and evaporated to afford **23-2** as a white solid. TLC Rf = 0.40 (25% EtOAc/Hexane)

^1H NMR (300 MHz, CDCl_3) δ 10.22 (s, 1H), 9.05 (d, $J = 1.8$ Hz, 1H), 8.27 (d, $J = 1.7$ Hz, 1H) 8.08 (d, $J = 1.8$ Hz, 1H), 7.10 (d, $J = 1.7$ Hz, 1H).

3-Amino-3-(furo[2,3-b]pyridin-5-yl)-propionic acid ethyl ester (23-3)

- 5 A solution containing aldehyde 23-2 (1.5 g, 10 mmol), ethyl hydrogen malonate (1.6 g, 20 mmol), and ammonium acetate (3.8 g, 50 mmol) in anhydrous ethanol (125 mL) was heated at reflux for 8 h. After cooling to room temperature, the solvent was evaporated and the residue partitioned between sat. sodium bicarbonate and EtOAc, the organic
10 layer removed, dried, and concentrated. Chromatography of the residue afforded the amino ester 23-3 as a waxy solid.
TLC $R_f = 0.5$ (20% 20:1:1 EtOH/ NH_4OH / H_2O - 80% EtOAc)
 ^1H NMR (300 MHz, CD_3OD) δ 8.34 (d, $J = 1.7$ Hz, 1H), 8.04 (d, $J = 1.7$ Hz, 1H), 7.72 (d, $J = 1.7$ Hz, 1H), 6.78 (d, $J = 1.7$ Hz, 1H), 4.62 (m, 1H), 4.13 (q,
15 $J = 7.5$ Hz, 2H), 3.20 (br, s, 2H), 2.76 (m, 2H), 1.23 (t, $J = 7.5$ Hz, 3H).


3-Furo[2,3-b]pyridin-6-yl-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (23-4)

- The title compound 23-4 was prepared from 2-9A and 23-3
20 using the procedure described in Scheme 2.
TLC $R_f = 0.30$ (50% 20:1:1 EtOH/ NH_4OH / H_2O - 50% EtOAc).
FAB MS Obs'd 450.1 (M+H).

3-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (23-5)

- 25 A solution of 23-4 (360 mg, 0.80 mmol) in MeOH (10 mL) was treated with 10% Pd/C (100 mg) and stirred under a hydrogen atmosphere for 18h. The catalyst was removed by filtration through celite and the residue chromatographed (75% 20:1:1 EtOH/ NH_4OH / H_2O -
30 25% EtOAc) to afford 23-5 as a colorless glass.
TLC $R_f = 0.30$ (50% 20:1:1 EtOH/ NH_4OH / H_2O - 50% EtOAc)
FAB MS Obs'd 452.2 (M+H).

Scheme 24

Furo[3,2-b]pyridine-5-carboxaldehyde (24-2)

- A solution of alcohol 24-1 (J.M. Hoffman, Jr., US Patent No. 4,808,595) was treated with excess MnO_2 (10 eq) and the mixture stirred at room temperature for 16 h, then filtered through Celite and evaporated to afford 24-2 as a white solid.
- ^1H NMR (300 MHz, CDCl_3) δ 10.18 (s, 1H), 8.92 (d, $J=1.8$ Hz, 1H), 8.17 (d, $J=1.7$ Hz, 1H) 7.89 (d, $J=1.8$ Hz, 1H), 7.10 (d, $J=1.7$ Hz, 1H).

3-Amino-3-(furo[3,2-b]pyridin-5-yl)-propionic acid ethyl ester (24-3)

A solution containing aldehyde 24-2 (1.5 g, 10 mmol), ethyl hydrogen malonate (1.6 g, 20 mmol), and ammonium acetate (3.8 g, 50 mmol) in anhydrous ethanol (125 mL) was heated at reflux for 8 h. After cooling to room temperature, the solvent was evaporated and the residue partitioned between sat. sodium bicarbonate and EtOAc, the organic layer removed, dried, and concentrated. Chromatography of the residue afforded the amino ester 24-3 as a waxy solid.

TLC R_f = 0.5 (20% 20:1:1 EtOH/NH₄OH/H₂O - 80% EtOAc)

¹H NMR (300 MHz, CD₃OD) δ 8.58 (d, J=1.7 Hz, 1H), 7.89 (d, J=1.7 Hz, 1H), 7.85 (d, J=1.7 Hz, 1H), 6.98 (d, J=1.7 Hz, 1H), 4.62 (t, J=7.2 Hz, 1H), 4.09 (q, J=7.5 Hz, 2H), 2.76 (m, 2H), 2.20 (br, s, 2H), 1.21 (t, J=7.5 Hz, 3H).

3-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (24-4)

The title compound 24-4 was prepared from 2-9A and 24-3 using the procedure described in Scheme 2.

TLC R_f = 0.56 (75% 20:1:1 EtOH/NH₄OH/H₂O - 25% EtOAc)

High resolution MS Calc'd.=450.2117, Obs'd = 450.2136.


3-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (24-5)

A solution of 24-4 (181 mg, 0.38 mmol) in acetic acid (5 mL) was treated with PtO₂ (100 mg) and stirred under a hydrogen atmosphere for 1h. The catalyst was removed by filtration through celite and the residue chromatographed (75% 20:1:1 EtOH/NH₄OH/H₂O - 25% EtOAc) to afford 24-5 as a colorless glass.

TLC R_f = 0.50 (75% 20:1:1 EtOH/NH₄OH/H₂O - 25% EtOAc)

High resolution MS Calc'd.=452.2298, Obs'd = 452.2238

Scheme 25



N-(S)-(2-Amino-phenyl)-3-tert-butoxycarbonylamino-succinamic acid methyl ester (25-3a)

- 5 A mixture of Boc-L-aspartic acid- β -methyl ester **25-1** (5.0g, 20.2 mmol), o-phenylenediamine **25-2a** (2.2 g, 20.2 mmol), EDC (3.9 g, 20.2 mmol), HOAT (0.28 g, 2.02 mmol), and NMM (6.7 mL, 60.7 mmol) in DMF (50 mL) was stirred for 18 h at ambient temperature. The solution was diluted with EtOAc (250 mL) and washed with sat. sodium
- 10 bicarbonate, water, and brine (50 mL each), then dried and evaporated to afford **25-3a** as a yellow solid.
- TLC R_f = 0.50 (95% CHCl_3 /5% isopropanol)
- ^1H NMR (300 MHz, CDCl_3) δ 8.10 (br,s, 1H), 7.23 (d, J = 7.8 Hz, 1H), 7.08 (t, J =7.8 Hz, 1H) 6.78 (m, 1H), 5.8 (br d, 1H), 4.65 (m, 1 H), 3.76 (s, 3H),
- 15 3.15 (dd, J =4.6, 16 Hz, 1H), 2.90 (dd, J = 5.1, 16 Hz, 1H), 1.48 (s, 9H).

3(S)-Amino-3-(benzimidazol-2-yl)-propionic acid methyl ester (25-4a)

Ester 25-3a (1.0 g, 3 mmol) was dissolved in acetic acid (50 mL) and heated to 65° for 2 h. The solvent was removed to afford the Boc-protected intermediate as a white solid. The crude material (920 mg, 2.43 mmol) was dissolved in EtOAc, cooled to 0°, and treated with HCl gas to give 25-4a as a tan solid.

¹H NMR (300 MHz, CD₃OD) δ 7.80 (m, 2H), 7.35 (m, 2H), 5.98 (m, 1H), 3.80 (m, 2H), 3.76 (s, 3H).

3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-1,8)naphthyridin-2-yl]propyl)imidazolidin-1-yl)propionic acid (25-5a)

The title compound 25-5a was prepared from 2-9A and 25-4a using the procedure described in Scheme 2.

TLC R_f = 0.30 (50% 20:1:1 EtOH/NH₄OH/H₂O-50% EtOAc).

FAB MS Obs'd 449.2 (M+H).


3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-1,8)naphthyridin-2-yl]propyl)imidazolidin-1-yl)propionic acid (25-5b)

The title compound 25-4b was prepared as described above substituting 3,4-diaminopyridine for o-phenylenediamine.

TLC R_f = 0.25 (50% 20:1:1 EtOH/NH₄OH/H₂O-50% EtOAc).

FAB MS Obs'd 450.2 (M+H).

Scheme 26



N-(S)-(2-Hydroxy-phenyl)-3-tert-butoxycarbonylamino-succinamic acid methyl ester (26-2)

A mixture of Boc-L-aspartic acid-β-methyl ester (25-1) (5.0g, 20.2 mmol), 2-amino phenol (26-1) (2.2 g, 20.2 mmol), EDC (3.9 g, 20.2 mmol), HOAT (0.28 g, 2.02 mmol), and NMM (6.7 mL, 60.7 mmol) in DMF (50 mL) was stirred for 18 h at ambient temperature. The solution was diluted with EtOAc (250 mL) and washed with sat. sodium bicarbonate, water, and brine (50 mL each), then dried, and evaporated and chromatographed on silica (EtOAc) to afford 26-2 as a white solid. TLC R_f = 0.55 (EtOAc)

¹H NMR (300 MHz, CDCl₃) δ 7.23 (d, J= 7.8 Hz, 1H), 6.89 (t, J=7.8 Hz, 1H), 6.78 (m, 1H), 5.68 (br d, 1H), 4.65 (m, 1 H), 3.76 (s, 3H), 3.15 (dd, J=4.6, 16 Hz, 1H), 2.90 (dd, J= 5.1, 16 Hz, 1H), 1.48 (s, 9H).

3(S)-Amino-(3-benzoxazol-2-yl)-propionic acid methyl ester (26-3)

Ester 26-2 (2.0 g, 6.0 mmol) was dissolved in anhydrous THF (150 mL) along with Ph₃P (1.58 g, 6.0 mmol). The resulting solution was cooled to 0°, and a solution of diethyl azodicarboxylate (1.53 g, 6.2 mmol) in THF (25 mL) was added dropwise. The cooling bath was removed and the solution stirred overnight at ambient temperature. The solution was concentrated and the residue chromatographed (75% EtOAc/Hexane) to afford the Boc-protected ester as a colorless glass. The crude material (1.8 g, 5.0 mmol) was dissolved in EtOAc, cooled to 0° and treated with HCl gas to give 26-3 as a tan solid.

¹H NMR (300 MHz, CD₃OD) δ 7.81 (m, 2H), 7.40 (m, 2H), 5.05 (t, J = 7.4 Hz, 1H), 3.72 (s, 3H), 3.30 (m, 2H).


3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid (26-4)

The title compound 26-4 was prepared from 2-9A and 26-3 using the procedure described in Scheme 2.

TLC R_f = 0.40 (50% 20:1:1 EtOH/NH₄OH/H₂O-50% EtOAc).

FAB MS Obs'd 450.3 (M+H).

SCHEME 27



1-Methyl-4-bromopyrazole (27-2)

Methyl iodide (8.47 mL, 136 mmol) was added to a mixture of 4-bromopyrazole 27-1 (10 g, 38 mmol), and K₂CO₃ (18.9 g, 136 mmol) in CH₃CN (150 mL) and the mixture stirred at room temperature for 16 h, then filtered and evaporated to yield 27-2 as a yellow oil.
¹H NMR (300 MHz, CDCl₃) δ 7.44(s, 1H), 7.38 (s, 1H), 3.90 (s, 3H).

3 (S)-Amino-3-(1-methyl-1H-pyrazol-4-yl)-propionic acid ethyl ester (27-3)

The bromide 27-2 was converted to the amino ester 27-3 following the procedure depicted in Scheme 1.

¹H NMR (300 MHz, CD₃OD) δ 7.81 (s, 1H), 7.58 (s, 1H), 4.80 (m, 1H), 4.05 (q, J = 7.0 Hz, 2 H), 3.89 (s, 3H), 3.00 (m, 2 H), 1.24 (t, J = 7.0 Hz, 3 H).


3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid (27-4)


The title compound 27-4 was prepared from 2-9A and 27-3 using the procedure described in Scheme 2.


TLC R_f = 0.24 (15:10:1:1 ethyl acetate/EtOH/water/NH₄OH).


¹H NMR (300 MHz, CD₃OD) δ 7.58 (s, 1H), 7.52 (d, J=7.3 Hz, 1H), 7.38 (s, 1H), 6.62 (d, J=7.3 Hz, 1H), 5.38 (m, 1H), 3.83 (s, 3H), 3.14-3.53 (9H), 2.97 (m, 2H), 2.80 (t, J=6.1 Hz, 2H), 2.67 (t, J=7.3 Hz, 2H), 1.93 (m, 4H).

SCHEME A
SYNTHESIS OF RADIOLIGAND FOR SPA ASSAY



SCHEME A (CONTINUED)

SCHEME A. cont'd



N-(4-Iodo-phenylsulfonylamino)-L-asparagine (A-2)

- To a stirred solution of acid **A-1** (4.39 g, 33.2 mmol), NaOH (1.49 g, 37.2 mmol), dioxane (30 ml) and H₂O (30 ml) at 0°C was added
- 5 pipsyl chloride (10.34 g, 34.2 mmol). After ~5 minutes, NaOH (1.49, 37.2 mmol), dissolved in 15 ml H₂O, was added followed by the removal of the cooling bath. After 2.0 h, the reaction mixture was concentrated. The residue was dissolved in H₂O (300 ml) and then washed with EtOAc.
- The aqueous portion was cooled to 0°C and then acidified with
- 10 concentrated HCl. The solid was collected and then washed with Et₂O to provide acid **A-2** as a white solid.
- ¹H NMR (300 MHz, D₂O) δ 7.86 (d, 2H, J=8Hz), 7.48 (d, 2H, J=8Hz) 3.70 (m, 1H), 2.39 (m, 2H).

15 2(S)-(4-Iodo-phenylsulfonylamino)-β-alanine (A-3)

- To a stirred solution of NaOH (7.14 g, 181.8 mmol) and H₂O (40 ml) at 0°C was added Br₂ (1.30 ml, 24.9 mmol) dropwise over a ten minute period. After ~5 minutes, acid **A-2** (9.9 g, 24.9 mmol), NaOH (2.00 g, 49.8 mmol) and H₂O (35 ml) were combined, cooled to 0°C and
- 20 then added in a single portion to the reaction. After stirring for

20 minutes at 0°C, the reaction was heated to 90°C for 30 minutes and then recooled to 0°C. The pH was adjusted to ~7 by dropwise addition of concentrated HCl. The solid was collected, washed with EtOAc, and then dried *in vacuo* to provide acid A-3 as a white solid.

5 ¹H NMR (300 MHz, D₂O) δ 8.02 (d, 2H, J=8Hz), 7.63 (d, 2H, J=8Hz), 4.36 (m, 1H), 3.51 (dd, 1H, J=5Hz, 13Hz) 3.21 (m, 1H).

Ethyl 2(S)-(4-iodo-phenylsulfonfylamino)-β-alanine-hydrochloride (A-4)

HCl gas was rapidly bubbled through a suspension of acid
10 A-3 (4.0 g, 10.81 mmol) in EtOH (50 ml) at 0°C for 10 minutes. The cooling bath was removed and the reaction was heated to 60°C. After 18 h, the reaction was concentrated to provide ester A-4 as a white solid.
¹H NMR (300 MHz, CD₃OD) δ 7.98 (d, 2H, J=8Hz), 7.63 (d, 2H, J=8Hz),
4.25 (q, 1H, J=5Hz), 3.92 (m, 2H), 3.33 (m, 1H), 3.06 (m, 1H), 1.01 (t, 3H,
15 J=7Hz).

Ethyl 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoate (A-5a)

A mixture of ester A-5 (700 mg, 2.63 mmol), (for
preparation, see: Scheme 29 of PCT International Application
20 Publication No. WO 95/32710, published December 7, 1995) 10% Pd/C (350 mg) and EtOH were stirred under 1 atm H₂. After 20 h, the reaction was filtered through a celite pad and then concentrated to provide ester A-5a as a brown oil.
TLC R_f = 0.23 (silica, 40% EtOAc/hexanes)
25 ¹H NMR (300 MHz, CDCl₃) δ 7.95 (d, 2H, J=8Hz), 7.26 (m, 3H), 6.43 (d, 1H, J=7Hz), 6.35 (d, 1H, J=8Hz), 4.37 (m, 4H), 3.05 (m, 2H), 2.91 (m, 2H), 1.39 (t, 3H, J=7Hz).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoic acid hydrochloride (A-6)

30 A suspension of ester A-5a (625 mg, 2.31 mmol) in 6N HCl (12 ml) was heated to 60°C. After ~20 h, the reaction was concentrated to give acid A-6 as a tan solid.
¹H NMR (300 MHz, CD₃OD) δ 7.96 (d, 2H, J=8Hz), 7.80 (m, 1H), 7.33 (d, 2H, J=8Hz), 6.84 (d, 1H, J=9Hz), 6.69 (d, 1H, J=7Hz), 3.09 (m, 4H).

35

Ethyl 4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-iodophenylsulfonylamino)-β-alanine (A-7)

- A solution of acid 15-6 (400 mg, 1.43 mmol), amine A-4 (686 mg, 1.57 mmol), EDC (358 mg, 1.86 mmol), HOBT (252 mg, 1.86 mmol), NMM (632 μl, 5.72 mmol) in DMF (10 ml) was stirred for ~20 h. The reaction was diluted with EtOAc and then washed with sat. NaHCO₃, brine, dried (MgSO₄) and concentrated. Flash chromatography (silica, EtOAc then 5% isopropanol/EtOAc) provided amide A-7 as a white solid.
- 10 TLC R_f = 0.4 (silica, 10% isopropanol/EtOAc)
¹H NMR (300 MHz, CD₃OD) δ 7.79 (d, 2H, J=9Hz) 7.61 (d, 2H, J=8Hz), 7.52 (d, 2H, J=9Hz), 7.29 (m, 1H), 7.27 (d, 2H, J=8Hz), 4.20 (m, 1H), 3.95 (q, 2H, J=7Hz), 3.66 (dd; 1H, J=6Hz, 14Hz), 3.49 (dd, 1H, J=8Hz, 13Hz), 3.01 (m, 2H), 2.86 (m, 2H), 1.08 (t, 3H, J=7Hz).

15

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-iodophenylsulfonylamino)-β-alanine (A-8)

- A solution of ester A-7 (200 mg, 0.3213 mmol) and 6N HCl (30 ml) was heated to 60°C. After ~20 h, the reaction mixture was concentrated. Flash chromatography (silica, 20:20:1:1 EtOAc/EtOH/NH₄OH/H₂O) provided acid A-8 as a white solid.
- 20 TLC R_f = 0.45 (silica, 20:20:1:1 EtOAc/EtOH/NH₄OH/H₂O)
¹H NMR (400 MHz, DMSO-d₆) δ 8.40 (m, 1H), 8.14 (Bs, 1H), 7.81 (d, 2H, J=8Hz), 7.62 (d, 2H, J=8Hz), 7.48 (d, 2H, J=8Hz), 7.27 (m, 3H), 6.34 (d, 1H, J=7Hz), 6.25 (d, 1H, J=8Hz), 5.85 (bs, 2H), 3.89 (bs, 1H), 3.35 (m, 2H), 2.97 (m, 2H), 2.79 (m, 2H).
- 25

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-(4-trimethylstannylphenylsulfonylamino)-β-alanine (A-9)

- 30 A solution of iodide A-8 (70 mg, 0.1178 mmol), [(CH₃)₃Sn]₂ (49 μl, 0.2356 mmol), Pd(PPh₃)₄ (5 mg) and dioxane (7 ml) was heated to 90°C. After 2 h, the reaction was concentrated and then purified by preparative HPLC (Delta-Pak C₁₈ 15 μM 100Å, 40 x 100 mm; 95:5 then 5:95 H₂O/CH₃CN) to provide the trifluoroacetate salt. The salt was

suspended in H₂O (10 ml), treated with NH₄OH (5 drops) and then lyophilized to provide amide A-9 as a white solid.

¹H NMR (400 MHz, DMSO-d₆) δ 8.40 (m, 1H), 8.18 (d, 1H, J=8Hz), 7.67 (m, 5H), 7.56 (d, 2H, J=8Hz), 7.29 (d, 2H, J=8Hz), 6.95-7.52 (m, 2H), 6.45 (bs, 2H), 4.00 (m, 1H), 3.50 (m, 1H), 3.33 (m, 1H), 2.97 (m, 2H), 2.86 (m, 2H).

4-[2-(2-Aminopyridin-6-yl)ethyl]benzoyl-2(S)-4-¹²⁵iodo-phenylsulfonfylamino-β-alanine (A-10)

10 An iodobead (Pierce) was added to a shipping vial of 5 mCi of Na¹²⁵I (Amersham, IMS30) and stirred for five minutes at room temperature. A solution of 0.1 mg of A-9 in 0.05 mL of 10% H₂SO₄/MeOH was made and immediately added to the Na¹²⁵I/iodobead vial. After stirring for three minutes at room temperature, approximately 0.04-0.05
15 mL of NH₄OH was added so the reaction mixture was at pH 6-7. The entire reaction mixture was injected onto the HPLC for purification [Vydac peptide-protein C-18 column, 4.6 x 250 mm, linear gradient of 10% acetonitrile (0.1% TFA):H₂O (0.1% TFA) to 90% acetonitrile (0.1% TFA):H₂O (0.1% TFA) over 30 minutes, 1 mL/min]. The retention time
20 of A-10 is 17 minutes under these conditions. Fractions containing the majority of the radioactivity were pooled, lyophilized and diluted with ethanol to give approximately 1 mCi of A-10, which coeluted on HPLC analysis with an authentic sample of A-8.

25 Instrumentation: Analytical and preparative HPLC was carried out using a Waters 600E Powerline Multi Solvent Delivery System with 0.1 mL heads with a Rheodyne 7125 injector and a Waters 990 Photodiode Array Detector with a Gilson FC203 Microfraction collector. For analytical and preparative HPLC, a Vydac peptide-protein
30 C-18 column, 4.6 x 250 mm was used with a C-18 Brownlee modular guard column. The acetonitrile used for the HPLC analyses was Fisher Optima grade. The HPLC radiodetector used was a Beckman 170 Radioisotope detector. A Vydac C-18 protein and peptide column, 3.9 x 250 mm was used for analytical and preparative HPLC. Solutions of

radioactivity were concentrated using a Speedvac vacuum centrifuge. Calibration curves and chemical concentrations were determined using a Hewlett Packard Model 8452A UV/Vis Diode Array Spectrophotometer. Sample radioactivities were determined in a Packard A5530 gamma
5 counter.

The test procedures employed to measure $\alpha\beta 3$ and $\alpha\beta 5$ binding and the bone resorption inhibiting activity of the compounds of the present invention are described below.

10 BONE RESORPTION-PIT ASSAY

When osteoclasts engage in bone resorption, they can cause the formation of pits in the surface of bone that they are acting upon. Therefore, when testing compounds for their ability to inhibit osteoclasts, it is useful to measure the ability of osteoclasts to excavate
15 these resorption pits when the inhibiting compound is present.

Consecutive 200 micron thick cross sections from a 6 mm cylinder of bovine femur diaphysis are cut with a low speed diamond saw (Isomet, Beuler, Ltd., Lake Bluff, IL). Bone slices are pooled, placed in a 10% ethanol solution and refrigerated until further use.

20 Prior to experimentation, bovine bone slices are ultrasonicated twice, 20 minutes each in H₂O. Cleaned slices are placed in 96 well plates such that two control lanes and one lane for each drug dosage are available. Each lane represents either triplicate or quadruplicate cultures. The bone slices in 96 well plates are sterilized by
25 UV irradiation. Prior to incubation with osteoclasts, the bone slices are hydrated by the addition of 0.1 ml α MEM, pH 6.9 containing 5% fetal bovine serum and 1% penicillin/streptomycin.

Long bones from 7-14 day old rabbits (New Zealand White Hare) are dissected, cleaned of soft tissue and placed in α MEM
30 containing 20 mM HEPES. The bones are minced using scissors until the pieces are <1 mm and transferred to a 50 ml tube in a volume of 25 ml. The tube is rocked gently by hand for 60 cycles, the tissue is sedimented for 1 min., and the supernatant is removed. Another 25 ml of medium is added to the tissue and rocked again. The second
35 supernatant is combined with the first. The number of cells is counted

excluding erythrocytes (typically $\sim 2 \times 10^7$ cells/ml). A cell suspension consisting of 5×10^6 /ml in α MEM containing 5% fetal bovine serum, 10 nM $1,25(\text{OH})_2\text{D}_3$, and penicillin-streptomycin is prepared. 200 ml aliquots are added to bovine bone slices (200 mm x 6 mm) and incubated
5 for 2 hrs. at 37°C in a humidified 5% CO_2 atmosphere. The medium is removed gently with a micropipettor and fresh medium containing test compounds is added. The cultures are incubated for 48 hrs., and assayed for c-telopeptide (fragments of the $\alpha 1$ chain of type I collagen) by Crosslaps for culture media (Herlev, Denmark).

10 Bovine bone slices are exposed to osteoclasts for 20-24 hrs and are processed for staining. Tissue culture media is removed from each bone slice. Each well is washed with 200 ml of H_2O , and the bone slices are then fixed for 20 minutes in 2.5% glutaraldehyde, 0.1 M cacodylate, pH 7.4. After fixation, any remaining cellular debris is
15 removed by 2 min. ultrasonication in the presence of 0.25 M NH_4OH followed by 2 X 15 min ultrasonication in H_2O . The bone slices are immediately stained for 6-8 min with filtered 1% toluidine blue and 1% borax.

After the bone slices have dried, resorption pits are counted
20 in test and control slices. Resorption pits are viewed in a Microphot Fx (Nikon) fluorescence microscope using a polarizing Nikon IGS filter cube. Test dosage results are compared with controls and resulting IC_{50} values are determined for each compound tested.

The appropriateness of extrapolating data from this assay to
25 mammalian (including human) disease states is supported by the teaching found in Sato, M., et al., Journal of Bone and Mineral Research, Vol. 5, No. 1, pp.31-40, 1990, which is incorporated by reference herein in its entirety. This article teaches that certain bisphosphonates have been used clinically and appear to be effective in
30 the treatment of Paget's disease, hypercalcemia of malignancy, osteolytic lesions produced by bone metastases, and bone loss due to immobilization or sex hormone deficiency. These same bisphosphonates are then tested in the resorption pit assay described above to confirm a

correlation between their known utility and positive performance in the assay.

EIB ASSAY

- 5 Duong *et al.*, *J. Bone Miner. Res.*, 8: S378 (1993) describes a system for expressing the human integrin $\alpha v \beta 3$. It has been suggested that the integrin stimulates attachment of osteoclasts to bone matrix, since antibodies against the integrin, or RGD-containing molecules, such as echistatin (European Publication 382 451), can effectively block
- 10 bone resorption.

Reaction Mixture:

1. 175 μ l TBS buffer (50 mM Tris•HCl pH 7.2, 150 mM NaCl, 1% BSA, 1 mM CaCl_2 , 1 mM MgCl_2).
- 15 2. 25 μ l cell extract (dilute with 100 mM octylglucoside buffer to give 2000 cpm/25 μ l).
3. ^{125}I -echistatin (25 μ l/50,000 cpm) (see EP 382 451).
4. 25 μ l buffer (total binding) or unlabeled echistatin (non-specific binding).

20

The reaction mixture was then incubated for 1 h at room temp. The unbound and the bound $\alpha v \beta 3$ were separated by filtration using a Skatron Cell Harvester. The filters (prewet in 1.5% polyethyleneimine for 10 mins) were then washed with the wash buffer (50

25 mM Tris HCl, 1mM $\text{CaCl}_2/\text{MgCl}_2$, pH 7.2). The filter was then counted in a gamma counter.

SPA ASSAY

30 MATERIALS:

1. Wheat germ agglutinin Scintillation Proximity Beads (SPA): Amersham
2. Octylglucopyranoside: Calbiochem
- 35 3. HEPES: Calbiochem

4. NaCl: Fisher
5. CaCl₂: Fisher
6. MgCl₂: SIGMA
7. Phenylmethylsulfonylfluoride (PMSF): SIGMA
- 5 8. Optiplate: PACKARD
9. Compound A-10 (specific activity 500-1000 Ci/mmol)
10. test compound
11. Purified integrin receptor: $\alpha_v\beta_3$ was purified from 293 cells overexpressing $\alpha_v\beta_3$ (Duong *et al.*, *J. Bone Min. Res.*, 8:S378, 1993) according to Pytela (*Methods in Enzymology*, 144:475, 1987)
12. Binding buffer: 50 mM HEPES, pH 7.8, 100 mM NaCl, 1 mM Ca²⁺/Mg²⁺, 0.5 mM PMSF
13. 50 mM octylglucoside in binding buffer: 50-OG buffer

PROCEDURE:

1. Pretreatment of SPA beads:
20 500 mg of lyophilized SPA beads were first washed four times with 200 ml of 50-OG buffer and once with 100 ml of binding buffer, and then resuspended in 12.5 ml of binding buffer.
2. Preparation of SPA beads and receptor mixture
25 In each assay tube, 2.5 μ l (40 mg/ml) of pretreated beads were suspended in 97.5 μ l of binding buffer and 20 μ l of 50-OG buffer. 5 μ l (~30 ng/ μ l) of purified receptor was added to the beads in suspension with stirring at room temperature for 30 minutes. The mixture was then centrifuged at 2,500 rpm in a
30 Beckman GPR Benchtop centrifuge for 10 minutes at 4°C. The pellets were then resuspended in 50 μ l of binding buffer and 25 μ l of 50-OG buffer.

3. Reaction

The following were sequentially added into Optiplate in corresponding wells:

- (i) Receptor/beads mixture (75 μ l)
- 5 (ii) 25 μ l of each of the following: compound to be tested, binding buffer for total binding or A-8 for non-specific binding (final concentration 1 μ M)
- (iii) A-10 in binding buffer (25 μ l, final concentration 40 pM)
- 10 (iv) Binding buffer (125 μ l)
- (v) Each plate was sealed with plate sealer from PACKARD and incubated overnight with rocking at 4°C

15 4. Plates were counted using PACKARD TOPCOUNT

5. % inhibition was calculated as follows:

A = total counts

B = nonspecific counts

20 C = sample counts

$$\% \text{ inhibition} = [(A-B)-(C-B)]/(A-B) \times 100$$

OCFORM ASSAY

Osteoblast-like cells (1.8 cells), originally derived from
25 mouse calvaria, were plated in CORNING 24 well tissue culture plates in α MEM medium containing ribo- and deoxyribonucleosides, 10% fetal bovine serum and penicillin-streptomycin. Cells were seeded at 40,000/well in the morning. In the afternoon, bone marrow cells were prepared from six week old male Balb/C mice as follows:

30 Mice were sacrificed, tibiae removed and placed in the above medium. The ends were cut off and the marrow was flushed out of the cavity into a tube with a 1 mL syringe with a 27.5 gauge needle. The marrow was suspended by pipetting up and down. The suspension was passed through >100 μ m nylon cell strainer. The resulting suspension
35 was centrifuged at 350 x g for seven minutes. The pellet was

resuspended, and a sample was diluted in 2% acetic acid to lyse the red cells. The remaining cells were counted in a hemacytometer. The cells were pelleted and resuspended at 1×10^6 cells/mL. 50 μ L was added to each well of 1.8 cells to yield 50,000 cells/well and 1,25-dihydroxy-vitamin D₃ (D₃) was added to each well to a final concentration of 10 nM. The cultures were incubated at 37°C in a humidified, 5% CO₂ atmosphere. After 48 h, the medium was changed. 72 h after the addition of bone marrow, test compounds were added with fresh medium containing D₃ to quadruplicate wells. Compounds were added again after 48 h with fresh medium containing D₃. After an additional 48 h., the medium was removed, cells were fixed with 10% formaldehyde in phosphate-buffered saline for 10 minutes at room temperature, followed by a 1-2 minute treatment with ethanol:acetone (1:1) and air dried. The cells were then stained for tartrate resistant acid phosphatase as follows:

The cells were stained for 10-15 minutes at room temperature with 50 mM acetate buffer, pH 5.0 containing 30 mM sodium tartrate, 0.3 mg/mL Fast Red Violet LB Salt and 0.1 mg/mL Naphthol AS -MX phosphate. After staining, the plates were washed extensively with deionized water and air dried. The number of multinucleated, positive staining cells was counted in each well.

α v β 5 ATTACHMENT ASSAY

Duong *et al.*, *J. Bone Miner. Res.*, 11: S290 (1996), describes a system for expressing the human α v β 5 integrin receptor.

Materials:

1. Media and solutions used in this assay are purchased from BRL/Gibco, except BSA and the chemicals are from Sigma.
2. Attachment medium: HBSS with 1 mg/ml heat-inactivated fatty acid free BSA and 2 mM CaCl₂.
3. Glucosaminidase substrate solution: 3.75 mM p-nitrophenyl N-acetyl-beta-D-glucosaminide, 0.1 M sodium citrate, 0.25% Triton, pH 5.0.

4. Glycine-EDTA developing solution: 50 mM glycine, 5 mM EDTA, pH 10.5.

Methods:

- 5 1. Plates (96 well, Nunc Maxi Sorp) were coated overnight at 4°C with human vitronectin (3 ug/ml) in 50 mM carbonate buffer (pH 9.6), using 100 µl/well. Plates were then washed 2X with DPBS and blocked with 2% BSA in DPBS for 2h at room temperature. After additional washes (2X) with
10 DPBS, plates were used for cell attachment assay.
2. 293 (αvβ5) cells were grown in MEM media in presence of 10% fetal calf serum to 90% confluence. Cells were then lifted from dishes with 1X Trypsin/EDTA and washed 3X with serum free MEM. Cells were resuspended in
15 attachment medium (3 X 10⁵ cells/ml).
3. Test compounds were prepared as a series of dilutions at 2X concentrations and added as 50 µl/well. Cell suspension was then added as 50 µl/well. Plates were incubated at 37°C with 55 CO₂ for 1 hour to allow attachment.
- 20 4. Non-adherent cells were removed by gently washing the plates (3X) with DPBS and then incubated with glucosaminidase substrate solution (100 µl/well), overnight at room temperature in the dark. To quantitate cell numbers, standard curve of glucosaminidase activity was
25 determined for each experiment by adding samples of cell suspension directly to wells containing the enzyme substrate solution.
5. The next day, the reaction was developed by addition of 185 µl/well of glycine/EDTA solution and reading absorbance at
30 405 nm using a Molecular Devices V-Max plate reader. Average test absorbance values (4 wells per test samples) were calculated. Then, the number of attached cells at each drug concentration was quantitated versus the standard curve of cells using the Softmax program.

35

EXAMPLE OF A PHARMACEUTICAL FORMULATION

As a specific embodiment of an oral composition, 100 mg of a compound of the present invention are formulated with sufficient
5 finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.





Representative compounds of the present invention were tested and found to bind to human $\alpha v \beta 3$ integrin. These compounds are generally found to have IC₅₀ values less than about 100 nM in the SPA
10 assay.



Representative compounds of the present invention were tested and generally found to inhibit $\geq 50\%$ the attachment of $\alpha v \beta 5$ expressing cells to plates coated with vitronectin at concentrations of about 1 μ M.

15 While the invention has been described and illustrated in reference to certain preferred embodiments thereof, those skilled in the art will appreciate that various changes, modifications and substitutions can be made therein without departing from the spirit and scope of the invention. For example, effective dosages other than the preferred doses
20 as set forth hereinabove may be applicable as a consequence of variations in the responsiveness of the mammal being treated for severity of bone disorders caused by resorption, or for other indications for the compounds of the invention indicated above. Likewise, the specific pharmacological responses observed may vary according to and
25 depending upon the particular active compound selected or whether there are present pharmaceutical carriers, as well as the type of formulation and mode of administration employed, and such expected variations or differences in the results are contemplated in accordance with the objects and practices of the present invention. It is intended,
30 therefore, that the invention be limited only by the scope of the claims which follow and that such claims be interpreted as broadly as is reasonable.


WHAT IS CLAIMED IS:

1. A compound having a structural formula selected from the group consisting of





, and




5

;

wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are substituted only with R¹⁰ and R¹²;

10

X is selected from the group consisting of



15

a 5- or 6-membered monocyclic aromatic or nonaromatic ring system having 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R¹ substituent and the ring

carbon atoms are unsubstituted or substituted with one or two R^1 substituents, and

5 a 9- to 14-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R^1 substituent and the ring carbon atoms are unsubstituted or substituted with one or two R^1 substituents;

10

Y is selected from the group consisting of

$-(CH_2)_m-$,
 $-(CH_2)_m-O-(CH_2)_n-$,
 $-(CH_2)_m-NR^4-(CH_2)_n-$,
 15 $-(CH_2)_m-S-(CH_2)_n-$,
 $-(CH_2)_m-SO-(CH_2)_n-$,
 $-(CH_2)_m-SO_2-(CH_2)_n-$,
 $-(CH_2)_m-O-(CH_2)_n-O-(CH_2)_p-$,
 $-(CH_2)_m-O-(CH_2)_n-NR^4-(CH_2)_p-$,
 20 $-(CH_2)_m-NR^4-(CH_2)_n-NR^4-(CH_2)_p-$,
 $-(CH_2)_m-O-(CH_2)_n-S-(CH_2)_p-$,
 $-(CH_2)_m-S-(CH_2)_n-S-(CH_2)_p-$,
 $-(CH_2)_m-NR^4-(CH_2)_n-S-(CH_2)_p-$,
 $-(CH_2)_m-NR^4-(CH_2)_n-O-(CH_2)_p-$,
 25 $-(CH_2)_m-S-(CH_2)_n-O-(CH_2)_p-$,
 $-(CH_2)_m-S-(CH_2)_n-NR^4-(CH_2)_p-$, and
 $-(CH_2)_m-Z-(CH_2)_n-$,

30 wherein Z is a 3- to 10-membered monocyclic or polycyclic aromatic or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R^1 substituent and the ring carbon atoms are unsubstituted or substituted with one or two R^1 substituents, and

wherein any methylene (CH₂) carbon atom in Y, other than in R⁴, can be substituted by one or two R³ substituents; and

- wherein R¹ and R² are each independently selected from the group
- 5 consisting of
- hydrogen, halogen, C₁₋₁₀ alkyl, C₃₋₈ cycloalkyl,
 - C₃₋₈ cycloheteroalkyl, C₃₋₈ cycloalkyl C₁₋₆ alkyl,
 - C₃₋₈ cycloheteroalkyl C₁₋₆ alkyl, aryl, aryl C₁₋₈ alkyl, amino,
 - 10 amino C₁₋₈ alkyl, C₁₋₃ acylamino, C₁₋₃ acylamino C₁₋₈ alkyl,
 - (C₁₋₆ alkyl)_pamino, (C₁₋₆ alkyl)_pamino C₁₋₈ alkyl,
 - C₁₋₄ alkoxy, C₁₋₄ alkoxy C₁₋₆ alkyl, hydroxycarbonyl,
 - hydroxycarbonyl C₁₋₆ alkyl, C₁₋₃ alkoxycarbonyl,
 - C₁₋₃ alkoxycarbonyl C₁₋₆ alkyl, hydroxycarbonyl-
 - 15 C₁₋₆ alkyloxy, hydroxy, hydroxy C₁₋₆ alkyl, C₁₋₆ alkyloxy-
 - C₁₋₆ alkyl, nitro, cyano, trifluoromethyl, trifluoromethoxy,
 - trifluoroethoxy, C₁₋₈ alkyl-S(O)_p, (C₁₋₈ alkyl)_paminocarbonyl,
 - C₁₋₈ alkyloxycarbonylamino, (C₁₋₈ alkyl)_paminocarbonyloxy,
 - (aryl C₁₋₈ alkyl)_pamino, (aryl)_pamino, aryl C₁₋₈
 - alkylsulfonylamino, and C₁₋₈ alkylsulfonylamino;
 - 20 or two R¹ substituents, when on the same carbon atom, are taken
 - together with the carbon atom to which they are attached to
 - form a carbonyl group;

- each R³ is independently selected from the group consisting of
- 25 hydrogen,
- aryl,
- C₁₋₁₀ alkyl,
- aryl-(CH₂)_r-O-(CH₂)_s-,
- aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
- 30 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
- aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
- aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
- aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
- halogen,

hydroxyl,
 oxo,
 trifluoromethyl,
 C₁₋₈ alkylcarbonylamino,
 5 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxy carbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 10 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 aryl C₁₋₅-alkylaminocarbonyl,
 aminocarbonyl,
 15 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 20 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 25 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 30 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 (aryl)_pamino,
 35 (aryl)_pamino C₁₋₆ alkyl,

- (aryl C₁₋₆ alkyl)pamino,
 (aryl C₁₋₆ alkyl)pamino C₁₋₆ alkyl,
 arylcarbonyloxy,
 aryl C₁₋₆ alkylcarbonyloxy,
 5 (C₁₋₆ alkyl)paminocarbonyloxy,
 C₁₋₈ alkylsulfonylamino,
 arylsulfonylamino,
 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
 arylsulfonylamino C₁₋₆ alkyl,
 10 aryl C₁₋₆ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
 C₁₋₈ alkoxycarbonylamino,
 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 aryloxy carbonylamino C₁₋₈ alkyl,
 15 aryl C₁₋₈ alkoxycarbonylamino,
 aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
 C₁₋₈ alkylcarbonylamino,
 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
 arylcarbonylamino C₁₋₆ alkyl,
 20 aryl C₁₋₆ alkylcarbonylamino,
 aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
 aminocarbonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)paminocarbonylamino,
 (C₁₋₈ alkyl)paminocarbonylamino C₁₋₆ alkyl,
 25 (aryl)paminocarbonylamino C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)paminocarbonylamino,
 (aryl C₁₋₈ alkyl)paminocarbonylamino C₁₋₆ alkyl,
 aminosulfonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)paminosulfonylamino,
 30 (C₁₋₈ alkyl)paminosulfonylamino C₁₋₆ alkyl,
 (aryl)paminosulfonylamino C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)paminosulfonylamino,
 (aryl C₁₋₈ alkyl)paminosulfonylamino C₁₋₆ alkyl,
 C₁₋₆ alkylsulfonyl,

5 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
arylsulfonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonyl,
aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
C₁₋₆ alkylcarbonyl,
C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
arylcarbonyl C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonyl,
aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
10 C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino,
aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
15 (C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
(aryl)_paminocarbonyl C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonyl, and
(aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl;
or two R³ substituents, when on the same carbon atom, are taken
20 together with the carbon atom to which they are attached to
form a carbonyl group or a cyclopropyl group,
wherein any of the alkyl groups of R³ are either unsubstituted or
substituted with one to three R¹ substituents, provided that each R³ is
selected such that in the resultant compound the carbon atom or atoms
25 to which R³ is attached is itself attached to no more than one
heteroatom;

each R⁴ is independently selected from the group consisting of
hydrogen,
30 aryl,
aminocarbonyl,
C₃₋₈ cycloalkyl,
amino C₁₋₆ alkyl,
(aryl)_paminocarbonyl,

(aryl C₁₋₅ alkyl)_paminocarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 C₁₋₈ alkyl,
 aryl C₁₋₆ alkyl,
 5 (C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
 C₁₋₈ alkylsulfonyl,
 C₁₋₈ alkoxycarbonyl,
 aryloxycarbonyl,
 10 aryl C₁₋₈ alkoxycarbonyl,
 C₁₋₈ alkylcarbonyl,
 arylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 15 aminosulfonyl,
 C₁₋₈ alkylaminosulfonyl,
 (aryl)_paminosulfonyl,
 (aryl C₁₋₈ alkyl)_paminosulfonyl,
 arylsulfonyl,
 20 arylC₁₋₆ alkylsulfonyl,
 C₁₋₆ alkylthiocarbonyl,
 arylthiocarbonyl, and
 aryl C₁₋₆ alkylthiocarbonyl,

wherein any of the alkyl groups of R⁴ are either unsubstituted or
 25 substituted with one to three R¹ substituents;

R⁵ and R⁶ are each independently selected from the group consisting of

hydrogen,
 C₁₋₁₀ alkyl,
 30 aryl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,

- aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 halogen,
 hydroxyl,
 5 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 10 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 15 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 20 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 25 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 30 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,

(aryl)_pamino,
(aryl)_pamino C₁₋₆ alkyl,
(aryl C₁₋₆ alkyl)_pamino,
(aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
5 arylcarbonyloxy,
aryl C₁₋₆ alkylcarbonyloxy,
(C₁₋₆ alkyl)_paminocarbonyloxy,
C₁₋₈ alkylsulfonylamino,
arylsulfonylamino,
10 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxy-carbonylamino,
15 C₁₋₈ alkoxy-carbonylamino C₁₋₈ alkyl,
aryloxy-carbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxy-carbonylamino,
aryl C₁₋₈ alkoxy-carbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino,
20 C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
25 (C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
30 aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,

- (aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
 C₁₋₆ alkylsulfonyl,
 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 arylsulfonyl C₁₋₆ alkyl,
 5 aryl C₁₋₆ alkylsulfonyl,
 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 C₁₋₆ alkylcarbonyl,
 C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 arylcarbonyl C₁₋₆ alkyl,
 10 aryl C₁₋₆ alkylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 C₁₋₆ alkylthiocarbonylamino,
 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 arylthiocarbonylamino C₁₋₆ alkyl,
 15 aryl C₁₋₆ alkylthiocarbonylamino,
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminocarbonyl, and
 20 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl;
 or R⁵ and R⁶ are taken together with the carbon atom to which they are
 attached to form a carbonyl group,
 wherein any of the alkyl groups of R⁵ or R⁶ are either unsubstituted or
 substituted with one to three R¹ substituents,
 25 and provided that each R⁵ and R⁶ are selected such that in the resultant
 compound the carbon atom to which R⁵ and R⁶ are attached is itself
 attached to no more than one heteroatom;
 R⁷ and R⁸ are each independently selected from the group consisting of
 30 hydrogen,
 C₁₋₁₀ alkyl,
 aryl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,

- aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 5 halogen,
 hydroxyl,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 10 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 C₃₋₈ cycloalkyl,
 (C₁₋₆ alkyl)_pamino,
 amino C₁₋₆ alkyl,
 15 arylaminocarbonyl,
 aryl C₁₋₅ alkylaminocarbonyl,
 aminocarbonyl,
 aminocarbonyl C₁₋₆ alkyl,
 hydroxycarbonyl,
 20 hydroxycarbonyl C₁₋₆ alkyl,
 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 25 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 30 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₆ alkoxy,
 aryl C₁₋₆ alkoxy,

- aryl C₁₋₆ alkyl,
(C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
(aryl)_pamino,
(aryl)_pamino C₁₋₆ alkyl,
5 (aryl C₁₋₆ alkyl)_pamino,
(aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
arylcarbonyloxy,
aryl C₁₋₆ alkylcarbonyloxy,
(C₁₋₆ alkyl)_paminocarbonyloxy,
10 C₁₋₈ alkylsulfonylamino,
arylcarbonylamino,
arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
15 aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
20 aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
25 aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
arylaminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino,
(C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
30 (aryl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,

- (C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
 (aryl)_paminosulfonylamino C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminosulfonylamino,
 (aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
 5 C₁₋₆ alkylsulfonyl,
 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 arylsulfonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkylsulfonyl,
 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 10 C₁₋₆ alkylcarbonyl,
 C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 arylcarbonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 15 C₁₋₆ alkylthiocarbonylamino,
 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 arylthiocarbonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylthiocarbonylamino,
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 20 (C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl C₁₋₈ alkyl)_paminocarbonyl,
 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl, and
 C₇₋₂₀ polycyclyl C₀₋₈ alkylsulfonylamino;
 25 wherein any of the alkyl groups of R⁷ and R⁸ are either unsubstituted or
 substituted with one to three R¹ substituents,
 and provided that each R⁷ and R⁸ are selected such that in the resultant
 compound the carbon atom to which R⁷ and R⁸ are attached is itself
 attached to no more than one heteroatom;
 30 R⁹ is selected from the group consisting of
 hydrogen,
 C₁₋₈ alkyl,
 aryl,

aryl C₁₋₈ alkyl,
 C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
 aryl C₁₋₈ alkylcarbonyloxy C₁₋₄ alkyl,
 C₁₋₈ alkylaminocarbonylmethylene, and
 5 C₁₋₈ dialkylaminocarbonylmethylene;

R₁₀, R₁₁, R₁₂ and R₁₃ are each independently selected from the group consisting of

10 hydrogen,
 C₁₋₈ alkyl,
 aryl,
 halogen,
 hydroxyl,
 aminocarbonyl,
 15 C₃₋₈ cycloalkyl,
 amino C₁₋₆ alkyl,
 (aryl)_paminocarbonyl,
 hydroxycarbonyl,
 (aryl C₁₋₅ alkyl)_paminocarbonyl,
 20 hydroxycarbonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkyl,
 (C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
 (aryl C₁₋₆ alkyl)_pamino C₂₋₆ alkyl,
 C₁₋₈ alkylsulfonyl,
 25 C₁₋₈ alkoxycarbonyl,
 aryloxycarbonyl,
 aryl C₁₋₈ alkoxycarbonyl,
 C₁₋₈ alkylcarbonyl,
 arylcarbonyl,
 30 aryl C₁₋₆ alkylcarbonyl,
 (C₁₋₈ alkyl)_paminocarbonyl,
 aminosulfonyl,
 C₁₋₈ alkylaminosulfonyl,
 (aryl)_paminosulfonyl,

- (aryl C₁₋₈ alkyl)_paminosulfonyl,
 C₁₋₆ alkylsulfonyl,
 arylsulfonyl,
 aryl C₁₋₆ alkylsulfonyl,
 5 aryl C₁₋₆ alkylcarbonyl,
 C₁₋₆ alkylthiocarbonyl,
 arylthiocarbonyl,
 aryl C₁₋₆ alkylthiocarbonyl,
 aryl-(CH₂)_r-O-(CH₂)_s-,
 10 aryl-(CH₂)_rS(O)_p-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-C(O)-N(R⁴)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-C(O)-(CH₂)_s-,
 aryl-(CH₂)_r-N(R⁴)-(CH₂)_s-,
 15 HC≡C-(CH₂)_t-,
 C₁₋₆ alkyl-C≡C-(CH₂)_t-,
 C₃₋₇ cycloalkyl-C≡C-(CH₂)_t-,
 aryl-C≡C-(CH₂)_t-,
 C₁₋₆ alkylaryl-C≡C-(CH₂)_t-,
 20 CH₂=CH-(CH₂)_t-,
 C₁₋₆ alkyl-CH=CH-(CH₂)_t-,
 C₃₋₇ cycloalkyl-CH=CH-(CH₂)_t-,
 aryl-CH=CH-(CH₂)_t-,
 C₁₋₆ alkylaryl-CH=CH-(CH₂)_t-,
 25 C₁₋₆ alkyl-SO₂-(CH₂)_t-,
 C₁₋₆ alkylaryl-SO₂-(CH₂)_t-,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₅ alkoxy,
 C₁₋₅ alkoxycarbonyl,
 30 (C₁₋₈ alkyl)_paminocarbonyl,
 C₁₋₆ alkylcarbonyloxy,
 (C₁₋₆ alkyl)_pamino,
 aminocarbonyl C₁₋₆ alkyl,
 C₁₋₆ alkoxy,

- aryl C₁₋₆ alkoxy,
(aryl)_pamino,
(aryl)_pamino C₁₋₆ alkyl,
(aryl C₁₋₆ alkyl)_pamino,
5 (aryl C₁₋₆ alkyl)_pamino C₁₋₆ alkyl,
arylcabonyloxy,
aryl C₁₋₆ alkylcabonyloxy,
(C₁₋₆ alkyl)_paminocabonyloxy,
C₁₋₈ alkylsulfonylamino,
10 arylsulfonylamino,
C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
15 C₁₋₈ alkoxycarbonylamino,
C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
20 C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
arylcabonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
25 aminocabonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocabonylamino,
(C₁₋₈ alkyl)_paminocabonylamino C₁₋₆ alkyl,
(aryl)_paminocabonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminocabonylamino,
30 (aryl C₁₋₈ alkyl)_paminocabonylamino C₁₋₆ alkyl,
aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,

(aryl C₁₋₈ alkyl)_paminosulfonylamino,
 (aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
 C₁₋₆ alkylsulfonyl,
 C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 5 arylsulfonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkylsulfonyl,
 aryl C₁₋₆ alkylsulfonyl C₁₋₆ alkyl,
 C₁₋₆ alkylcarbonyl,
 C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 10 arylcarbonyl C₁₋₆ alkyl,
 aryl C₁₋₆ alkylcarbonyl,
 aryl C₁₋₆ alkylcarbonyl C₁₋₆ alkyl,
 C₁₋₆ alkylthiocarbonylamino,
 C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 15 arylthiocarbonylamino C₁₋₆ alkyl,
 aryl C₁₋₆ alkylthiocarbonylamino,
 aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
 (C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl,
 (aryl)_paminocarbonyl C₁₋₆ alkyl,
 20 (aryl C₁₋₈ alkyl)_paminocarbonyl, and
 (aryl C₁₋₈ alkyl)_paminocarbonyl C₁₋₆ alkyl; or
 R¹⁰ and R¹² are taken together with the carbon atoms to which
 they are attached to form a 5- to 7-membered monocyclic aromatic
 or nonaromatic ring system having 0, 1, 2, 3, or 4 heteroatoms
 25 selected from the group consisting of N, O, and S wherein the ring
 nitrogen atoms are unsubstituted or substituted with one R¹
 substituent and the ring carbon atoms are unsubstituted or
 substituted with one or two R¹ substituents,
 and wherein any of the alkyl groups of R¹⁰, R¹¹, R¹², and R¹³ are either
 30 unsubstituted or substituted with one to three R¹ substituents;

wherein



each m is independently an integer from 0 to 6;

each n is independently an integer from 0 to 6


- each p is independently an integer from 0 to 2;
 each r is independently an integer from 1 to 3;
 each s is independently an integer from 0 to 3;
 each t is independently an integer from 0 to 3; and
 5 each v is independently an integer from 0 to 2;

and the pharmaceutically acceptable salts thereof.

2. The compound of Claim 1 having a structural
 10 formula selected from the group consisting of



and



15

wherein the dotted line a represents a single or a double bond, provided
 that when a represents a double bond, the double bond carbon atoms are
 substituted only with R¹⁰ and R¹²;


20

X is

a 6-membered monocyclic aromatic ring system having 1 or 2 nitrogen atoms wherein each ring carbon atom is unsubstituted or substituted with one R¹ substituent, or

- 5 a 9- to 14-membered polycyclic ring system, wherein one or more of the rings is aromatic, and wherein the polycyclic ring system has 0, 1, 2, 3 or 4 heteroatoms selected from the group consisting of N, O, and S wherein the ring nitrogen atoms are unsubstituted or substituted with one R¹ substituent and the ring carbon atoms are
- 10 unsubstituted or substituted with one or two R¹ substituents.

3. The compound of Claim 2 having structural formula




15


wherein the dotted line a represents a single or a double bond, provided that when a represents a double bond, the double bond carbon atoms are substituted only with R¹⁰ and R¹²; and

20

X is selected from the group consisting of




4. The compound of Claim 3 having structural formula



5

5. The compound of Claim 4 wherein X is



6. The compound of Claim 5 wherein Y is selected from the group consisting of

10

- (CH₂)_m-,
- (CH₂)_m-O-(CH₂)_n-,
- (CH₂)_m-NR⁴-(CH₂)_n-,
- (CH₂)_m-S-(CH₂)_n-,
- 15 -(CH₂)_m-SO-(CH₂)_n-,
- (CH₂)_m-SO₂-(CH₂)_n-,
- (CH₂)_m-O-(CH₂)_n-O-(CH₂)_p-,
- (CH₂)_m-O-(CH₂)_n-NR⁴-(CH₂)_p-,
- (CH₂)_m-NR⁴-(CH₂)_n-NR⁴-(CH₂)_p-, and
- 20 -(CH₂)_m-NR⁴-(CH₂)_n-O-(CH₂)_p-,

wherein any methylene (CH₂) carbon atom in Y, other than in R⁴, can be substituted by one or two R³ substituents.

25

7. The compound of Claim 6 wherein Y is selected from the group consisting of

(CH₂)_m, (CH₂)_m-S-(CH₂)_n, and (CH₂)_m-NR⁴-(CH₂)_n,
5 wherein any methylene (CH₂) carbon atom in Y, other than in R⁴, can be substituted by one or two R³ substituents,
m and n are integers from 0-4,
and v is 0.

10 8. The compound of Claim 7 wherein Y is

(CH₂)_m or (CH₂)_m-NR⁴-(CH₂)_n,
wherein any methylene (CH₂) group in Y, other than in R⁴, can be substituted by one or two R³ substituents.

15

9. The compound of Claim 8 wherein each R³ is independently selected from the group consisting of

hydrogen,
fluoro,
20 trifluoromethyl,
aryl,
C₁₋₈ alkyl,
arylC₁₋₆ alkyl
hydroxyl,
25 oxo,
arylaminocarbonyl,
aryl C₁₋₅ alkylaminocarbonyl,
aminocarbonyl, and
aminocarbonyl C₁₋₆ alkyl;

30

and each R⁴ is independently selected from the group consisting of

hydrogen,
aryl,
C₃₋₈ cycloalkyl,
35 C₁₋₈ alkyl,

C₁₋₈ alkylcarbonyl,
arylcarbonyl,
C₁₋₆ alkylsulfonyl,
arylsulfonyl,
5 arylC₁₋₆alkylsulfonyl,
arylC₁₋₆alkylcarbonyl,
C₁₋₈alkylaminocarbonyl,
arylC₁₋₅alkylaminocarbonyl,
arylC₁₋₈alkoxycarbonyl, and
10 C₁₋₈alkoxycarbonyl.

10. The compound of Claim 9 wherein R⁶, R⁷, and R⁸ are each hydrogen and R⁵ is selected from the group consisting of
hydrogen,
15 aryl,
C₁₋₈ alkyl,
aryl-C≡C-(CH₂)_t-,
aryl C₁₋₆ alkyl,
CH₂=CH-(CH₂)_t-, and
20 HC≡C-(CH₂)_t-.

11. The compound of Claim 10 wherein R¹⁰, R¹¹, R¹², and R¹³ are each independently selected from the group consisting of hydrogen, aryl, C₁₋₆alkyl, and arylC₁₋₆alkyl.

25

12. The compound of Claim 10 wherein R⁹ is selected from the group consisting of hydrogen, methyl, and ethyl.

13. The compound of Claim 12 wherein R⁹ is hydrogen.
30

14. The compound of Claim 9 wherein R⁵, R⁶, and R⁸ are each hydrogen and R⁷ is selected from the group consisting of
hydrogen,
aryl,

- C₁₋₈ alkylcarbonylamino,
C₁₋₈ alkylsulfonylamino,
arylcarbonylamino,
arylsulfonylamino,
5 C₁₋₈ alkylsulfonylamino C₁₋₆ alkyl,
arylsulfonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylsulfonylamino,
aryl C₁₋₆ alkylsulfonylamino C₁₋₆ alkyl,
C₁₋₈ alkoxycarbonylamino,
10 C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
aryloxycarbonylamino C₁₋₈ alkyl,
aryl C₁₋₈ alkoxycarbonylamino,
aryl C₁₋₈ alkoxycarbonylamino C₁₋₈ alkyl,
C₁₋₈ alkylcarbonylamino C₁₋₆ alkyl,
15 arylcarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylcarbonylamino,
aryl C₁₋₆ alkylcarbonylamino C₁₋₆ alkyl,
aminocarbonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminocarbonylamino,
20 (C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
(aryl)_paminocarbonylamino C₁₋₆ alkyl,
arylaminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino,
(aryl C₁₋₈ alkyl)_paminocarbonylamino C₁₋₆ alkyl,
25 aminosulfonylamino C₁₋₆ alkyl,
(C₁₋₈ alkyl)_paminosulfonylamino,
(C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl)_paminosulfonylamino C₁₋₆ alkyl,
(aryl C₁₋₈ alkyl)_paminosulfonylamino,
30 (aryl C₁₋₈ alkyl)_paminosulfonylamino C₁₋₆ alkyl,
C₁₋₆ alkylthiocarbonylamino,
C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl,
arylthiocarbonylamino C₁₋₆ alkyl,
aryl C₁₋₆ alkylthiocarbonylamino, and

aryl C₁₋₆ alkylthiocarbonylamino C₁₋₆ alkyl.

15. The compound of Claim 14 wherein R⁷ is selected from the group consisting of

- 5 hydrogen,
 aryl,
 C₁₋₈ alkylcarbonylamino,
 aryl C₁₋₆ alkylcarbonylamino,
 arylcarbonylamino,
10 C₁₋₈ alkylsulfonylamino,
 aryl C₁₋₆ alkylsulfonylamino,
 arylsulfonylamino,
 C₁₋₈ alkoxy carbonylamino,
 aryl C₁₋₈ alkoxy carbonylamino,
15 arylaminocarbonylamino,
 (C₁₋₈ alkyl)_paminocarbonylamino,
 (aryl C₁₋₈ alkyl)_paminocarbonylamino,
 (C₁₋₈ alkyl)_paminosulfonylamino, and
 (aryl C₁₋₈ alkyl)_paminosulfonylamino.

20

16. The compound of Claim 15 wherein R¹⁰, R¹¹, R¹², and R¹³ are each independently selected from the group consisting of hydrogen, aryl, C₁₋₆alkyl, and arylC₁₋₆ alkyl.

25 17. The compound of Claim 15 wherein R⁹ is selected from the group consisting of hydrogen, methyl, and ethyl.

 18. The compound of Claim 17 wherein R⁹ is
 hydrogen.

30

19. The compound of Claim 9 selected from the group consisting of

- 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 3(R)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3(S)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 35

3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

5 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

10

3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,

15 3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,

3-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl}-propionic acid,

20 3(S)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3(R)-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

25

3-(6-Methoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

30 3(S)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3(R)-(6-Ethoxypyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
trifluoroacetate salt,

5 3(S)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
bis(trifluoroacetate) salt,

3(R)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
10 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

15 3(S)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

20

3-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3-(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3- [3-
25 (5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-
propionic acid,

3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3- [3-
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-
30 propionic acid,

3-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3- [3-
(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-
propionic acid,

35

- 3(S)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
5 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(6-Methylamino-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 15 3-(2-Fluoro-biphenyl-4-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
20 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 25 3-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 30 3(R)-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(4-Ethoxy-3-fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
35 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 5 3(R)-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(5-Ethoxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 10 3(S)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(R)-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
15 [1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3-(5-Hydroxy-pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 20 2(S)-Benzenesulfonylamino-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,
- 3(S)-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-
imidazolidin-1-yl}-pent-4-enoic acid,
- 25 3(R)-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-
imidazolidin-1-yl}-pent-4-enoic acid,
- 3-{2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-
30 imidazolidin-1-yl}-pent-4-enoic acid,
- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-{3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-
yl]-propyl]-2-oxo-imidazolidin-1-yl}-propionic acid,

3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,

5 3-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,

3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid,

10 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(R)-(5-ethoxy-pyridin-3-yl)-propionic acid,

3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3-(5-ethoxy-pyridin-3-yl)-propionic acid,

15

3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,

20 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,

25 3-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,

30 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,

35 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl}propionic acid,

- 3-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15 3-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 3(R)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 15 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 20 3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 25 3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 35 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

5 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

10

3-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

15 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

20 3-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

25

3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

30 3-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

20. The compound of Claim 19 selected from the group consisting of

5 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3(R)-(2,3-Dihydro-benzofuran-6-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

10 3(S)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

15 3(R)-(3-Fluorophenyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3(S)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

20 3(R)-(Quinolin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3(S)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

25 3(R)-(Ethynyl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

30 3(S)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

3(R)-(Pyridin-3-yl)-3-{2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl}-propionic acid,

- 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,
- 3(R)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-4-methyl-imidazolidin-1-yl)-propionic acid,
- 5 3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 10 3(R)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 15 3(R)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 20 bis(trifluoroacetate) salt,
- 3(R)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3(S)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(R)-(6-Amino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 30 3(S)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 35

3-(R)-(4-Methyl-3-oxo-3,4-dihydro-2H-benzo[1,4]oxazin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 5 3(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

10

3(S)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 15 3(R)-(2-Fluoro-biphenyl-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 20 3(R)-(2-Oxo-2,3-dihydro-benzoxazol-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

25

3(R)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 30 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

3(R)-(5-Ethoxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,

- 3(S)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 5 3(R)-(5-Hydroxy-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 10 3(R)-(2-Oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-pent-4-enoic acid,
- 3(S)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 15 3(R)-(5-Ethoxy-pyridin-3-yl)-3-(3-[3-[6-(4-methoxy-benzylamino)-pyridin-2-yl]-propyl]-2-oxo-imidazolidin-1-yl)-propionic acid,
- 20 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(S)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 3-[3-[3-(6-Amino-pyridin-2-yl)-propyl]-2-oxo-imidazolidin-1-yl]-3(R)-(5-ethoxy-pyridin-3-yl)-propionic acid,
- 25 3(S)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 30 3(R)-(2-Oxo-2,3-dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 35 3(S)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,

- 3(R)-(2,3-Dihydro-1H-4-oxa-1,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 5 3(S)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 10 3(R)-(3-Oxo-3,4-dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 15 3(S)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(R)-(3,4-Dihydro-2H-1-oxa-4,5-diaza-naphthalen-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 25 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(R)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 35 3(S)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 30 3(R)-(2,3-Dihydrofuro[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 35 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

- 3(R)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 5 3(S)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(2,3-Dihydrofuro[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 10 3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(R)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
15 [1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 20 3(R)-(1H-Imidazo[4,5-c]pyridin-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 25 3(R)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-
2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
30 [1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 3(R)-(1-Methyl-1H-pyrazol-4-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]-imidazolidin-1-yl)propionic acid,
- 35 and the pharmaceutically acceptable salts thereof.

21. The compound of Claim 20 selected from the group consisting of

- 5 3(S)-(2,3-Dihydro-benzofuran-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(Quinolin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 10 3(S)-(Pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 3(S)-(6-Methoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 15 3(S)-(6-Ethoxypyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 20 3(S)-(4-Methoxyquinolin-7-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid, bis(trifluoroacetate) salt,
- 3(S)-(6-Methylamino-pyridin-3-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 25 3(S)-(4-Ethoxy-3-fluorophenyl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)-propyl]-imidazolidin-1-yl)-propionic acid,
- 30 3(S)-(Furo[2,3-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 3(S)-(Furo[3,2-b]pyridin-6-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,
- 35

3(S)-(Benzimidazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-
[1,8]naphthyridin-2-yl)propyl]imidazolidin-1-yl)propionic acid,

3(S)-(Benzoxazol-2-yl)-3-(2-oxo-3-[3-(5,6,7,8-tetrahydro-[1,8]naphthyridin-
5 2-yl)propyl]imidazolidin-1-yl)propionic acid,

and the pharmaceutically acceptable salts thereof.

22. A pharmaceutical composition comprising a
10 compound according to Claim 1 and a pharmaceutically acceptable
carrier.

23. A pharmaceutical composition made by combining a
compound according to Claim 1 and a pharmaceutically acceptable
15 carrier.

24. A process for making a pharmaceutical composition
comprising combining a compound according to Claim 1 and a
pharmaceutically acceptable carrier.

20

25. The composition of Claim 22 which further comprises
an active ingredient selected from the group consisting of

- a) an organic bisphosphonate or a pharmaceutically
acceptable salt or ester thereof,
- 25 b) an estrogen receptor modulator,
- c) a cytotoxic/antiproliferative agent,
- d) a matrix metalloproteinase inhibitor,
- e) an inhibitor of epidermal-derived, fibroblast-derived, or
platelet-derived growth factors,
- 30 f) an inhibitor of VEGF,
- g) an inhibitor of Flk-1/KDR, Flt-1, Tck/Tie-2, or Tie-1,
- h) a cathepsin K inhibitor, and
- i) a prenylation inhibitor, such as a farnesyl transferase
inhibitor or a geranylgeranyl transferase inhibitor or a dual
35 farnesyl/geranylgeranyl transferase inhibitor;

and mixtures thereof.

26. The composition of Claim 25 wherein said active ingredient is selected from the group consisting of

- 5 a) an organic bisphosphonate or a pharmaceutically acceptable salt or ester thereof,
 - b) an estrogen receptor modulator, and
 - c) a cathepsin K inhibitor;
- and mixtures thereof.

10

27. The composition of Claim 26 wherein said organic bisphosphonate or pharmaceutically acceptable salt or ester thereof is alendronate monosodium trihydrate.

15

28. The composition of Claim 25 wherein said active ingredient is selected from the group consisting of

- a) a cytotoxic/antiproliferative agent,
 - b) a matrix metalloproteinase inhibitor,
 - c) an inhibitor of epidermal-derived, fibroblast-derived, or
 - 20 platelet-derived growth factors,
 - d) an inhibitor of VEGF, and
 - e) an inhibitor of Flk-1/KDR, Flt-1, Tck/Tie-2, or Tie-1;
- and mixtures thereof.

25

29. A method of eliciting an integrin receptor antagonizing effect in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound according to Claim 1.

30

30. The method of Claim 29 wherein the integrin receptor antagonizing effect is an $\alpha v \beta 3$ antagonizing effect.

31. The method of Claim 30 wherein the $\alpha v \beta 3$ antagonizing effect is selected from the group consisting of inhibition of

bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular degeneration, inflammation, viral disease, and tumor growth.

32. The method of Claim 31 wherein the $\alpha v \beta 3$
5 antagonizing effect is the inhibition of bone resorption.

33. The method of Claim 29 wherein the integrin receptor
antagonizing effect is an $\alpha v \beta 5$ antagonizing effect.

10 34. The method of Claim 33 wherein the $\alpha v \beta 5$
antagonizing effect is selected from the group consisting of inhibition of
restenosis, angiogenesis, diabetic retinopathy, macular degeneration,
inflammation, and tumor growth.

15 35. The method of Claim 29 wherein the integrin receptor
antagonizing effect is a dual $\alpha v \beta 3 / \alpha v \beta 5$ antagonizing effect.

36. The method of Claim 35 wherein the dual $\alpha v \beta 3 / \alpha v \beta 5$
antagonizing effect is selected from the group consisting of inhibition of
20 bone resorption, restenosis, angiogenesis, diabetic retinopathy, macular
degeneration, inflammation, viral disease, and tumor growth.

37. The method of Claim 29 wherein the integrin
antagonizing effect is an $\alpha v \beta 6$ antagonizing effect.
25

38. The method of Claim 37 wherein the $\alpha v \beta 6$
antagonizing effect is selected from the group consisting of
angiogenesis, inflammatory response, and wound healing.

30 39. A method of eliciting an integrin receptor
antagonizing effect in a mammal in need thereof, comprising
administering to the mammal a therapeutically effective amount of the
composition of Claim 22.

40. A method of treating or preventing a condition mediated by antagonism of an integrin receptor in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 22.

5

41. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 22.

10

42. A method of inhibiting bone resorption in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 26.

15

43. A method of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of the composition of Claim 28.

20

44. A method of treating tumor growth in a mammal in need thereof, comprising administering to the mammal a therapeutically effective amount of a compound according to Claim 1 in combination with radiation therapy.

INTERNATIONAL SEARCH REPORT

Internati	Application No
PCT/US 98/26568	

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 C07D401/06 C07D401/14 C07D519/00 A61K31/415
 A61K31/44 //(C07D519/00,498:00,471:00),(C07D519/00,491:00,
 471:00),(C07D519/00,471:00,471:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 051 829 A (A. NATTERMANN & CIE. GMBH) 19 May 1982 see claims 1,20; examples 1,18,48	1,22
A	WO 95 17397 A (MERCK & CO., INC.) 29 June 1995 cited in the application see claims 1,4,5	1,22
A	WO 95 32710 A (MERCK & CO., INC.) 7 December 1995 cited in the application see claims 11-14,17	1,22
A	WO 91 18888 A (G. D. SEARLE & CO.) 12 December 1991 see claims 1,22; examples 7,9,10,24,27	1,22
	-/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

7 April 1999

Date of mailing of the international search report

13/04/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax (+31-70) 340-3016

Authorized officer

Hass, C

INTERNATIONAL SEARCH REPORT

Internat. J. Application No.
PCT/US 98/26568

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 93 16674 A (SMITHKLINE BEECHAM CORP.) 2 September 1993 see claims 26-28 -----	1,22
A	US 4 460 595 A (D. R. ADAMS ET AL.) 17 July 1984 see claim 1; tables 3,6 -----	1,22
A	EP 0 006 718 A (BEECHAM GROUP PLC) 9 January 1980 see claims 1-3,11 -----	1,22

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/26568

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 29-44
because they relate to subject matter not required to be searched by this Authority, namely:
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Although claims 29-44 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Claims Nos.: 29-44

Rule 39.1(iv) PCT - Method for treatment of the human or animal body by therapy

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internal / Application No

PCT/US 98/26568

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 51829	A	19-05-1982	DE 3042466 A	16-06-1982
			AT 10626 T	15-12-1984
			DK 495781 A	12-05-1982
			GR 74703 A	05-07-1984
			IE 52079 B	10-06-1987
			JP 57108073 A	05-07-1982
			US 4349684 A	14-09-1982
			ZA 8107696 A	27-10-1982
WO 9517397	A	29-06-1995	US 5416099 A	16-05-1995
			AU 1441695 A	10-07-1995
WO 9532710	A	07-12-1995	AU 701776 B	04-02-1999
			AU 2586895 A	21-12-1995
			CA 2190870 A	07-12-1995
			EP 0760658 A	12-03-1997
			JP 10501222 T	03-02-1998
			US 5741796 A	21-04-1998
WO 9118888	A	12-12-1991	AU 7903091 A	31-12-1991
			CA 2083735 A	26-11-1991
			EP 0531382 A	17-03-1993
WO 9316674	A	02-09-1993	AU 3664593 A	13-09-1993
			AU 701541 B	28-01-1999
			AU 4186797 A	08-01-1998
			AU 703236 B	18-03-1999
			AU 6482698 A	16-07-1998
			CA 2129897 A	12-08-1993
			EP 0626969 A	07-12-1994
			US 5648373 A	15-07-1997
			US 5663053 A	02-09-1997
			CA 2061061 A	12-08-1993
US 4460595	A	17-07-1984	GB 1598073 A	16-09-1981
			AU 509364 B	08-05-1980
			AU 3161277 A	21-06-1979
			CA 1097660 A	17-03-1981
			CA 1101870 A	26-05-1981
			CH 638190 A	15-09-1983
			DE 2755727 A	29-06-1978
			DK 551677 A	19-06-1978
			FI 773813 A	19-06-1978
			FR 2374313 A	13-07-1978
			IE 46118 B	23-02-1983
			JP 53082781 A	21-07-1978
			NL 7713973 A	20-06-1978
			SE 7714372 A	19-06-1978
			US 4367338 A	04-01-1983
			BE 861957 A	16-06-1981
			ZA 7707336 A	25-10-1978
EP 6718	A	09-01-1980	AU 4806879 A	20-12-1979
			JP 55004385 A	12-01-1980